


Georgia Department of Natural Resources
American Shad Habitat Plan



Submitted to the Atlantic States Marine Fisheries Commission as a requirement of Amendment 3 to the Interstate Management Plan for Shad and River Herring

Approved February 6, 2014

Shad Habitat Plan-Georgia

Altamaha River

Habitat Assessment

The Altamaha River is formed by the confluence of the Ocmulgee and Oconee rivers and flows approximately 220 km before emptying into the Atlantic Ocean near Darien, GA. Including its longest tributary (the Ocmulgee River) the Altamaha River runs for approximately 756 km making it the seventh longest river in the U.S. that is entirely within one state. The Altamaha River drainage basin covers an area of approximately 36,000 km² with its headwaters arising near Atlanta, GA for the Ocmulgee River and near Athens, GA for the Oconee River. There are no dams directly on the Altamaha, though there are dams on both the Oconee and the Ocmulgee rivers.

With no barriers directly on the Altamaha all historical estuarine habitat remains available to juvenile and migrating adult shad.

Historical evidence suggests that American shad once occurred in the Altamaha Basin at least as far upstream as the vicinity of Covington, GA in the Ocmulgee River Basin and near the city of Athens, GA in the Oconee River Basin [Bryson 1826; Baird 1884; Bill Frazier, U. S. Fish and Wildlife Service (retired), 2001, personal communication; Elizabeth Reitz, University of Georgia, 2007, personal communication]. However, the construction of dams has limited the migrations. Most of these structures are still in place and continue to serve as barriers to nearly 6,000 acres of potential riverine shad habitat.

American shad currently occur from the mouth of the Altamaha River to the East Juliette Hydroelectric Dam on the Ocmulgee River (approximately river km 570) and Sinclair Dam on the Oconee River (approximately river km 446). Approximately 70% of the historical riverine habitat currently remains available to migrating adult American shad.

Threats Assessment

- 1. Migration Barriers-** Full utilization of all potential spawning habitat in the Altamaha River Basin could entail modification of at least nine dams in the Oconee Basin, seven dams in the Ocmulgee Basin, and one dam in the Ochopee Basin to facilitate fish passage.

Action 1: Develop a plan for establishing fish passage at barriers in the Altamaha River system.

Regulatory Agencies/Contacts: USFWS, NMFS, FERC, USACE, Georgia Department of Natural Resources (GA DNR), dam owners and operators, and federal and state legislators.

Goal/Target: Establish fish passage at all dams in the Altamaha basin, where passage is determined to be feasible.

Progress: GA DNR has developed an American shad restoration plan for the Altamaha River Basin which includes the implementation of fishways as a restoration strategy. The plan calls for utilizing Section 18 of the Federal Power Act, which provides the U.S. Departments of Commerce and Interior mandatory conditioning authority to prescribe fish passage during the Federal Energy Regulation Commission (FERC) licensing process for hydroelectric facilities. The FERC-licensed hydroelectric facilities in the Altamaha Basin that are within the historic range of the American shad should have fish passage provisions included in their upcoming licenses, when passage is determined to be feasible.

For FERC-licensed facilities that already have a spawning population directly below them (e.g., currently East Juliette Hydroelectric Dam, Sinclair Dam), fish passage should be evaluated and implemented as soon as feasible (or upon FERC relicensing). For all other FERC-licensed facilities, fish passage should be provided in a stepwise fashion upon the establishment of spawning runs directly below these structures (upon fish passage at all downstream structures).

For non-FERC-licensed dams resource agencies should work with owners to explore passage opportunities such as fishways, breaching, or removal. Where feasible, obsolete or non-functioning barriers to migration should be removed or breached.

East Juliette Hydroelectric Dam

A fish passage prescription for East Juliette Hydroelectric dam has been completed. However, negotiations between the Services and project operator are still ongoing and construction of the fishway has not been initiated.

Cost: Unknown

Timeline: Unknown

Action 2: Potentially conduct experimental trap and transport operations.

Regulatory Agencies/Contacts: Georgia Department of Natural Resources (GA DNR), ASMFC, USFWS, NMFS, FERC, USACE, dam owners and operators, and federal and state legislators.

Goal/Target: Assess of upstream migratory behavior and level of passage at partial barriers and to provide access to additional spawning habitat that may be more suitable than that available below downstream barriers.

Progress: Experimental trap and transport operations are listed as a potential method for assessing migratory behavior, partial barrier passage, and allow for potential spawning at previously unavailable habitat. GA DNR has no immediate plans to initiate trap and transport activities at this time.

Cost: Unknown

Timeline: Unknown

- 2. Dissolved Oxygen-**While there have not been any dissolved oxygen issues identified within the Altamaha River itself, segments of tributary rivers and streams have been identified as not having sufficient assimilative capacity to maintain dissolved oxygen levels of 5mg/L or greater at maximum permitted discharge levels under low flow conditions.

Action 1: Develop a regional water plan that recommends appropriate water management practices to ensure healthy aquatic ecosystems.

Regulatory Agencies/Contacts: Georgia Department of Natural Resources (GA DNR)- Environmental Protection Division (EPD), Wildlife Resources Division (WRD), and Coastal Resources Division (CRD), state legislators, and local municipalities

Goal/Target: Ensure water quantity remains adequate to support all life stages of American shad and other aquatic organisms in the Altamaha River.

Progress: In 2008, the Georgia General Assembly, as part of the Statewide Comprehensive Water Management Plan, established 10 regional water planning councils that encompassed the 14 major river systems within Georgia. With technical guidance from GA EPD, these councils were tasked with developing regional water plans that outlined management practices to meet future water needs for both water quantity and water quality through 2050. In November 2011, the ten regional water plans were officially adopted by GA EPD.

The Altamaha Council recommended a suite of surface water quality management practices in a phased approach to address water quality issues, including stream segments with limited localized dissolved oxygen assimilative capacity and insufficient wastewater permit capacity (GA EPD 2011a. These recommendations include such practices as the additional sustainable development of groundwater and surface water in areas with sufficient water supply; best management practices for water quality issues such as non-point source runoff, nutrient loadings, and TMDLs in the region; and additional educational and ordinance practices.

For the Altamaha Region, 75 impaired stream reaches (total impaired length of 915 miles) and 2 impaired lakes (total impaired area of 390 acres) have been identified. The majority of impairments are due to low dissolved oxygen and fecal coliform. Total maximum daily loads have been completed for 71 of the impaired stream reaches and for both of the impaired lakes.

Cost: Unknown

Timeline: Regional water plan extends through 2050

3. Competition and Predation by Invasive Species-Flathead catfish and blue catfish have been introduced into that Altamaha River system through unauthorized stockings. A significant portion of both flathead catfish and blue catfish diets are comprised of fish, and due to their large adult size (>60 lbs) they have the potential to consume both adult and juvenile American shad. Flathead catfish were first documented in the Ocmulgee River in the early-1970's and have now colonized the entire Altamaha River system. Abundance of flathead catfish rapidly expanded from approximately 1980 through the late-1990's. Electrofishing catch rates by weight peaked at 274 kg/hr in 1993 and by number at 108 fish/hr in 2004. Since 2000, electrofishing catch rates have ranged from 43-108 fish/hr. The average size of the flathead catfish in the Altamaha River peaked at approximately 3.5 kg in the mid-1990's and has since decreased to approximately 1 kg. A diet analysis of flathead catfish was completed during the months of June-September of 1997 and found the dominant prey items to be centrarchid spp. and ictalurid spp (Weller and Robbins, 2001). No *Alosa* spp. were identified in the stomach of flathead catfish during this study, but consumed juvenile American and/or hickory shad could have been unidentifiable due to extensive digestion.

Blue catfish were first detected in the Altamaha River in 2006 and their abundance has steadily increased. In 2011, blue catfish electrofishing CPUE was 29 fish/hr. It is expected that the abundance of this species will continue to increase for several more years. Stomach contents of 257 blue catfish were analyzed in the summer of 2010 and it was found that *Alosa* spp. comprised 0.4% by number of prey items consumed (Bonvechio et al. 2012). This majority of the blue catfish in this study were relatively small (59.5% < 300 mm) so as larger blue catfish become more abundant utilization of *Alosa* spp as a prey item may increase.

Action 1: Management of invasive catfish species.

Regulatory Agencies/Contacts: GA DNR

Progress: GA DNR completed experimental electrofishing removals of flathead catfish from the Altamaha River system during the 1990s in an effort to restore native fish redbreast sunfish and bullhead spp populations that had been adversely impacted. These efforts were discontinued due to the large nature of the river, budget reductions, and shifts in angler attitudes.

Cost: Unknown

Timeline: Discontinued

Ogeechee River

Habitat Assessment

The Ogeechee River originates in the Georgia piedmont and flows for approximately 425 km while crossing the fall line, sandhill region, and the coast plain before emptying into the Atlantic Ocean in Ossabaw Sound. The Ogeechee River watershed encompasses approximately 14,300 km². Tidal influence typically extends to rkm 72 and the fresh/saltwater interface occurs approximately 56 km upstream from the mouth of the river. No manmade barriers are present the entire length of the Ogeechee River so all historical riverine and estuarine habitats remain available to juvenile and migrating adult American shad.

Threats Assessment

- 1. Instream Flow-** The Georgia Environmental Protection Division (EPD) conducted resource assessments to predict resource conditions based on projection population growth and resulting water demands through 2050. Based on these predictions peak season agricultural irrigation may result in potential in-stream flow shortages in the Ogeechee Basin (GA EPD 2011b). The stream flow may fall below the in-stream flow target during summer low flow periods after meeting upstream irrigation needs.

Action 1: Develop a regional water plan that recommends appropriate water management practices to ensure healthy aquatic ecosystems.

Regulatory Agencies/Contacts: Georgia Department of Natural Resources (GA DNR)-Environmental Protection Division (EPD), Wildlife Resources Division (WRD), and Coastal Resources Division (CRD), USFWS, NMFS, FERC, US EPD, USACE, federal and state legislators, and local municipalities.

Goal/Target: Ensure water quantity remains adequate to support all life stages of American shad and other aquatic organisms in the Ogeechee River.

Progress: In 2008, the Georgia General Assembly, as part of the Statewide Comprehensive Water Management Plan, established 10 regional water planning councils that encompassed the 14 major river systems within Georgia. With technical guidance from GA EPD, these councils were tasked with developing regional water plans that outlined management practices to meet future water needs for both water quantity and water quality through 2050. In November 2011, the ten regional water plans were officially adopted by GA EPD.

To prevent potential shortages in meeting in-stream flow needs, the plan encompassing the Ogeechee River calls for more aggressive water conservation practices and development of drought management practices for the agricultural users/permittees in the Upper Ogeechee River Basin (GA EPD 2011b). The Council also recommends in-stream flow studies (to determine what flow levels are appropriate for protecting aquatic

life) and additional stream flow monitoring in the Ogeechee River Basin (to confirm the frequency and magnitude of predicted in-stream flow shortages).

Cost: Unknown

Timeline: Regional water plan extends through 2050

- 2. Point Source Discharges-** In May 2011, the Ogeechee River experienced a large-scale fish kill that affected multiple species including American shad. The upper extent of the kill was below the only industrial discharge above the kill area.

Action 1: Develop and implement permits and monitoring to avoid future fish kills.

Regulatory Agencies/Contacts: Georgia Department of Natural Resources (GA DNR)-Environmental Protection Division (EPD), Wildlife Resources Division (WRD), US EPD, and appropriate private industrial operators.

Goal/Target: Ensure water quality remains adequate to support all life stages of American shad and other aquatic organisms in the Ogeechee River.

Progress: After the 2011 fish kill, GA EPD reviewed and revised the existing discharge permit for King America Finishing in attempt to prevent future fish kills related to their discharge. GA EPD has since closely monitored water quality in this area of the Ogeechee River.

Cost: Unknown

Timeline: Currently ongoing

Satilla River

Habitat Assessment

The Satilla River originates in Ben Hill County near the town of Fitzgerald, GA and flows for approximately 378 km before emptying into the Atlantic Ocean in St. Andrews Sound. The Satilla River watershed encompasses approximately 10,000 km² of Georgia's coastal plain. Tidal influence typically extends to rkm 93 and the fresh/saltwater interface occurs approximately 32 km upstream from the mouth of the river. No manmade barriers are present the entire length of the Satilla River so all historical riverine and estuarine habitats remain available to juvenile and migrating adult American shad.

Threats Assessment

- 1. Competition and Predation by Invasive Species-**Flathead catfish were introduced into that Satilla River system through unauthorized stockings in the mid-1990s and blue catfish were collected by GA DNR in 2012. A significant portion of flathead catfish diets are comprised of fish, and due to their large adult size (>60 lbs) they have the potential to consume both adult and juvenile American shad.

Action 1: Management of invasive catfish species.

Regulatory Agencies/Contacts: GA DNR

Progress: GA DNR initiated electrofishing removals of flathead catfish from the Satilla River in 1996 with existing manpower and funding in an effort to preserve native fish species, specifically redbreast sunfish and bullhead spp. Flathead abundance continued to increase despite these efforts, which were limited due to manpower and fiscal limitations. Native fish populations were also showing early signs of decline. In 2006, Georgia's legislature appropriated funding for dedicated positions and equipment to conduct extensive flathead catfish removal efforts on the Satilla River. Since 2007, approximately 28,000 flathead catfish weighing over 68,000 lbs have been removed from the Satilla River. Over time, these efforts have resulted in a significant reduction in the flathead catfish biomass and appear to be preserving the abundance of native species.

Blue catfish abundance is extremely low, with only a few individual being collected in 2012 and none thus far in 2013. GA DNR suspects that these fish may have colonized the Satilla River from the Altamaha River via the intercostal water way during a high flow period, due to their relatively high tolerance to brackish water.

Cost: Unknown

Timeline: Ongoing

- 2. Dissolved Oxygen-** Dissolved oxygen levels below 3 mg/L occur during low flow events in the months of July-September in an approximately a 30 km segment of the tidally influenced portion of the Satilla River. The Satilla River naturally has a low assimilative capacity and resulting low DO levels during summer low flow periods, therefore it may not be possible to maintain DO levels above 3 mg/L at all times. However, the actions listed below will still be beneficial.

Action 1: Develop a TMDL implementation plan.

Regulatory Agencies/Contacts: Georgia Department of Natural Resources (GA DNR)- Environmental Protection Division (EPD), Wildlife Resources Division (WRD), and Coastal Resources Division (CRD), state legislators, and local municipalities

Goal/Target: Reduce organic loads to sustain acceptable DO levels.

Progress: GA DNR worked with representatives of local municipalities and conservation groups and developed a TMDL implementation plan that included a suite of management measure to reduce organic carbon, Total Nitrogen, and Total Phosphorous inputs in order to improve dissolved oxygen levels in the Satilla River.

Cost: Unknown

Timeline: Unknown

Action 2: Develop a regional water plan that recommends appropriate water management practices to ensure healthy aquatic ecosystems.

Regulatory Agencies/Contacts: Georgia Department of Natural Resources (GA DNR)-Environmental Protection Division (EPD), Wildlife Resources Division (WRD), and Coastal Resources Division (CRD), USFWS, NMFS, FERC, US EPD, USACE, federal and state legislators, and local municipalities.

Goal/Target: Ensure water quantity remains adequate to support all life stages of American shad and other aquatic organisms in the Satilla River.

Progress: In 2008, the Georgia General Assembly, as part of the Statewide Comprehensive Water Management Plan, established 10 regional water planning councils that encompassed the 14 major river systems within Georgia. With technical guidance from GA EPD, these councils were tasked with developing regional water plans that outlined management practices to meet future water needs for both water quantity and water quality through 2050. In November 2011, the ten regional water plans were officially adopted by GA EPD.

The Suwannee-Satilla-St Marys Council recommended a suite of surface water quality management practices in a phased approach to address water quality gaps, including stream segments with limited localized dissolved oxygen assimilative capacity and insufficient wastewater permit capacity (GA EPD 2011c). Specific actions to add/improve infrastructure and improve flow and water quality conditions were identified and recommended. These recommendations include such practices as the additional sustainable development of groundwater and surface water in areas with sufficient water supply; best management practices for water quality issues such as non-point source runoff, nutrient loadings, and TMDLs in the region; and additional educational and ordinance practices.

Cost: Unknown

Timeline: Regional water plan extends through 2050

3. **Instream Flow-** The Georgia Environmental Protection Division (EPD) conducted resource assessments on current and predicted resource conditions based on projected population growth and resulting water demands through 2050. These assessments concluded that instream flow shortages were present under current and future demands in portions of the Satilla Basin.

Action 1: Develop a regional water plan that recommends appropriate water management practices to ensure healthy aquatic ecosystems.

Regulatory Agencies/Contacts: Georgia Department of Natural Resources (GA DNR)-Environmental Protection Division (EPD), Wildlife Resources Division (WRD), and Coastal Resources Division (CRD), USFWS, NMFS, FERC, US EPD, USACE, federal and state legislators, and local municipalities.

Goal/Target: Ensure water quantity remains adequate to support all life stages of American shad and other aquatic organisms in the Satilla River.

Progress: The Satilla River water management plan was officially adopted by GA EPD in November 2011 and recommended a suite of management practices, including those that reduce net consumption, replace surface water use with groundwater use, and improve data on frequency and magnitude of gaps (GA EPD 2011c).

Cost: Unknown

Timeline: Regional water plan extends through 2050

St. Marys River

Habitat Assessment

The St. Marys River originates in the Okefenokee Swamp and flows for approximately 203 km before emptying into the Atlantic Ocean in Cumberland Sound while forming the eastern portion of the border between Florida and Georgia. The St. Marys watershed encompasses approximately 3,350 km² of which 59% is in Georgia and 41% in Florida. Tidal influence typically extends to rkm 88 and the fresh/saltwater interface occurs approximately 33 km upstream from the mouth of the river. No manmade barriers are present the entire length of the St. Marys River so all historical riverine and estuarine habitats remain available to juvenile and migrating adult American shad.

Threats Assessment

1. **Dissolved Oxygen-** Dissolved oxygen levels below 3 mg/L occur during low flow events in the months of July-September months of July-September in an approximately a

40 km segment of the tidally influenced portion of the St. Marys River. The St Marys River naturally has a low assimilative capacity and resulting low DO levels during summer low flow periods, therefore it may not be possible to maintain DO levels above 3 mg/L at all times. However, the actions listed below will still be beneficial.

Action 1: Develop a TMDL implementation plan.

Regulatory Agencies/Contacts: Georgia Department of Natural Resources (GA DNR)- Environmental Protection Division (EPD), Wildlife Resources Division (WRD), and Coastal Resources Division (CRD), FL FWC, FL DEP, St. Johns Water Management District, state legislators, and local municipalities

Goal/Target: Reduce organic loads to sustain acceptable DO levels.

Progress: GA DNR worked with representatives of local municipalities and conservation groups and developed a TMDL implementation plan that included a suite of management measure to reduce organic inputs in order to improve dissolved oxygen levels in the St. Marys River.

Cost: Unknown

Timeline: Unknown

Action 2: Develop a regional water plan that recommends appropriate water management practices to ensure healthy aquatic ecosystems.

Regulatory Agencies/Contacts: Georgia Department of Natural Resources (GA DNR)- Environmental Protection Division (EPD), Wildlife Resources Division (WRD), and Coastal Resources Division (CRD), USFWS, NMFS, FERC, US EPD, USACE, federal and state legislators, and local municipalities.

Goal/Target: Ensure water quantity remains adequate to support all life stages of American shad and other aquatic organisms in the St. Marys River.

Progress: In 2008, the Georgia General Assembly, as part of the Statewide Comprehensive Water Management Plan, established 10 regional water planning councils that encompassed the 14 major river systems within Georgia. With technical guidance from GA EPD, these councils were tasked with developing regional water plans that outlined management practices to meet future water needs for both water quantity and water quality through 2050. In November 2011, the ten regional water plans were officially adopted by GA EPD.

The Suwannee-Satilla-St Marys Council recommended a suite of surface water quality management practices in a phased approach to address water quality gaps, including stream segments with limited localized dissolved oxygen assimilative capacity and insufficient wastewater permit capacity (GA EPD 2011c). Specific actions to

add/improve infrastructure and improve flow and water quality conditions were identified and recommended. These recommendations include such practices as the additional sustainable development of groundwater and surface water in areas with sufficient water supply; best management practices for water quality issues such as non-point source runoff, nutrient loadings, and TMDLs in the region; and additional educational and ordinance practices.

Cost: Unknown

Timeline: Regional water plan extends through 2050

References

- Baird, S. 1884. The fisheries and fishery industries of the United States. United States Commission of Fish and Fisheries, Washington, D.C., Section I, Plate 214, page 598.
- Bonvechio et al. 2012. Diet and Population Metrics of the Introduced Blue Catfish (*Ictalurus furcatus*) on the Altamaha River, Georgia. Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies. 65 (2011):112-118.
- Bryson H. 1826. Travels of Henry Bryson.
[Http://www.arptaft.org/Henry%20Bryson.htm](http://www.arptaft.org/Henry%20Bryson.htm).
- Evans, J et. al. 2012. Priority restoration and management actions for the American shad in the Altamaha River Basin, Georgia. Georgia Department of Natural Resources. 2012. 29 pp.
- GA EPD. 2011a. Altamaha Regional Water Plan. 147pp.
[Http://www.georgiawaterplanning.org/documents/ALT_Adopted_RWP_000.pdf](http://www.georgiawaterplanning.org/documents/ALT_Adopted_RWP_000.pdf)
- GA EPD. 2011b. Savannah-Upper Ogeechee Regional Water Plan. 119 pp.
[Http://www.georgiawaterplanning.org/documents/SUO_Adopted_RWP_000.pdf](http://www.georgiawaterplanning.org/documents/SUO_Adopted_RWP_000.pdf)
- GA EDP. 2011c. Suwannee Satilla Regional Water Plan. 155 pp.
[Http://www.georgiawaterplanning.org/documents/SSA_Adopted_RWP_000.pdf](http://www.georgiawaterplanning.org/documents/SSA_Adopted_RWP_000.pdf)
- Hill, A. 2005. Diadromous fish restoration plan for the middle Savannah River: strategy and implementation schedule. U.S. Fish Wildlife Service National Marine Fisheries Service. 2005. 28 pp.
- Weller, R. R., and C. Robbins. 2001. Food habits of flathead catfish in the Altamaha River System, Georgia. Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies. 53(1999):35-41.