

# Research Priorities and Recommendations to Support Interjurisdictional Fisheries Management

## ATLANTIC STRIPED BASS

(Full Citation: Atlantic States Marine Fisheries Commission. 2013. Research Priorities and Recommendations to Support Interjurisdictional Fisheries Management. Special Report # 89. ASMFC, Arlington, VA. 58pp.)

### Fishery-Dependent Priorities

#### *Moderate*

- Develop studies to provide information on gear specific discard mortality rates and to determine the magnitude of bycatch mortality.<sup>1</sup>
- Improve estimates of striped bass harvest removals in coastal areas during wave 1 and inland waters of all jurisdictions year round.
- Evaluate the percentage of fishermen using circle hooks.<sup>2</sup>

### Fishery-Independent Priorities

#### *Moderate*

- Develop a refined and cost-efficient, fisheries-independent coastal population index for striped bass stocks.

### Modeling / Quantitative Priorities

#### *High*

- Develop a method to integrate catch-at-age and tagging models to produce a single estimate of F and stock status.<sup>3</sup>
- Develop a spatially and temporally explicit catch-at-age model incorporating tag based movement information.<sup>4</sup>
- Review model averaging approach to estimate annual fishing mortality with tag based models. Review validity and sensitivity to year groupings.<sup>5</sup>
- Develop methods for combining tag results from programs releasing fish from different areas on different dates.
- Examine potential biases associated with the number of tagged individuals, such as gear specific mortality (associated with trawls, pound nets, gill nets, and electrofishing), tag induced mortality, and tag loss.<sup>6</sup>

---

<sup>1</sup> Literature search and some modeling work completed.

<sup>2</sup> Work ongoing in New York through the Hudson River Angler Diary, Striped Bass Cooperative Angler Program, and ACCSP elogbook.

<sup>3</sup> Model developed, but the tagging data overwhelms the model. Issues remain with proper weighting.

<sup>4</sup> Model developed with Chesapeake Bay and the rest of the coast as two fleets. However, no tagging data has been used in the model.

<sup>5</sup> Work ongoing by Striped Bass Tagging Subcommittee to evaluate the best years to use for the IRCR and the periods to use for the MARK models.

- Develop field or modeling studies to aid in estimation of natural mortality or other factors affecting the tag return rate.

#### ***Moderate***

- Develop maturity ogives applicable to coastal migratory stocks.
- Examine methods to estimate annual variation in natural mortality.<sup>7</sup>
- Develop reliable estimates of poaching loss from striped bass fisheries.
- Improve methods for determining population sex ratio for use in estimates of SSB and biological reference points.
- Evaluate truncated matrices and covariate based tagging models.

#### ***Low***

- Examine issues with time saturated tagging models for the 18 inch length group.
- Develop tag based reference points.

### **Life History, Biological, and Habitat Priorities**

#### ***High***

- Continue in-depth analysis of migrations, stock compositions, etc. using mark-recapture data.<sup>8</sup>
- Continue evaluation of striped bass dietary needs and relation to health condition.<sup>9</sup>
- Continue analysis to determine linkages between the mycobacteriosis outbreak in Chesapeake Bay and sex ratio of Chesapeake spawning stock, Chesapeake juvenile production, and recruitment success into coastal fisheries.

#### ***Moderate***

- Examine causes of different tag based survival estimates among programs estimating similar segments of the population.
- Continue to conduct research to determine limiting factors affecting recruitment and possible density implications.
- Conduct study to calculate the emigration rates from producer areas now that population levels are high and conduct multi-year study to determine inter-annual variation in emigration rates.

#### ***Low***

- Determine inherent viability of eggs and larvae.
- Conduct additional research to determine the pathogenicity of the IPN virus isolated from striped bass to other warm water marine species, such as flounder, menhaden, shad, and largemouth bass.

#### ***Additional Habitat Research Recommendations***

- Passage facilities should be designed specifically for passing striped bass for optimum efficiency at passing this species.

---

<sup>6</sup> Gear specific survival being examined in Hudson River.

<sup>7</sup> Ongoing work by the Striped Bass Tagging Subcommittee

<sup>8</sup> Ongoing through Cooperative Winter Tagging Cruise and striped bass charter boat tagging trips. See Cooperative Winter Tagging Cruise 20 Year Report.

<sup>9</sup> Plans for a stomach content collection program in the Chesapeake Bay by the Chesapeake Bay Ecological Foundation.

- Conduct studies to determine whether passing migrating adults upstream earlier in the year in some rivers would increase striped bass production and larval survival, and opening downstream bypass facilities sooner would reduce mortality of early emigrants (both adult and early-hatched juveniles).
- All state and federal agencies responsible for reviewing impact statements and permit applications for projects or facilities proposed for striped bass spawning and nursery areas shall ensure that those projects will have no or only minimal impact on local stocks, especially natal rivers of stocks considered depressed or undergoing restoration.<sup>10</sup>
- Federal and state fishery management agencies should take steps to limit the introduction of compounds which are known to be accumulated in striped bass tissues and which pose a threat to human health or striped bass health.
- Every effort should be made to eliminate existing contaminants from striped bass habitats where a documented adverse impact occurs.
- Water quality criteria for striped bass spawning and nursery areas should be established, or existing criteria should be upgraded to levels that are sufficient to ensure successful striped bass reproduction.
- Each state should implement protection for the striped bass habitat within its jurisdiction to ensure the sustainability of that portion of the migratory stock. Such a program should include: inventory of historical habitats, identification of habitats presently used, specification of areas targeted for restoration, and imposition or encouragement of measures to retain or increase the quantity and quality of striped bass essential habitats.
- States in which striped bass spawning occurs should make every effort to declare striped bass spawning and nursery areas to be in need of special protection; such declaration should be accompanied by requirements of non-degradation of habitat quality, including minimization of non-point source runoff, prevention of significant increases in contaminant loadings, and prevention of the introduction of any new categories of contaminants into the area. For those agencies without water quality regulatory authority, protocols and schedules for providing input on water quality regulations to the responsible agency should be identified or created, to ensure that water quality needs of striped bass stocks are met.<sup>11</sup>
- ASMFC should designate important habitats for striped bass spawning and nursery areas as HAPC.
- Each state should survey existing literature and data to determine the historical extent of striped bass occurrence and use within its jurisdiction. An assessment should be conducted of those areas not presently used for which restoration is feasible.

## **Management, Law Enforcement, and Socioeconomic Priorities**

### ***Moderate***

- Examine the potential public health trade-offs between the continued reliance on the use of high minimum size limits (28 inches) on coastal recreational anglers and its long-term effects on enhanced PCB contamination among recreational stakeholders.<sup>12</sup>

---

<sup>10</sup> Ongoing in New York.

<sup>11</sup> Significant habitat designations completed in the Hudson River and New York Marine Districts.

<sup>12</sup> Samples collected from two size groups ( $\geq 28$  inches and 20-26 inches) in Pennsylvania and processed by the Department of Environmental Protection to compare contamination of the two size groups.