

2001 REVIEW OF THE  
ATLANTIC STATES MARINE FISHERIES COMMISSION  
FISHERY MANAGEMENT PLAN FOR  
**TAUTOG**  
*(Tautoga onitis)*

Prepared by

Heather Stirratt (ASMFC)

Tautog Plan Review Team

Paul Caruso (MA)  
David Simpson (CT)  
Frank Steimle (NMFS)  
Najih Lazar (ASMFC)

June 2001

**2001 REVIEW OF THE  
ASMFC FISHERY MANAGEMENT PLAN FOR  
TAUTOG (*Tautoga onitis*)**

**I. Status of Fishery Management Plan Development**

In May 1993 the ASMFC voted to develop a Fishery Management Plan for Tautog. The primary rationale was the vulnerability of tautog to overfishing. Additional concerns centered around localized overfishing and rising commercial fishing effort and landings. States declaring interest in tautog were: Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Delaware, Maryland, and Virginia. Plan development began in 1994 and the Tautog FMP was approved in March 1996. In May 1997 the Commission approved Addendum 1 to the Tautog FMP. Addendum 1 included *de minimus* specifications and adjustments to the compliance schedule. Required management measures began April 1, 1998. In November 1999 the Commission approved Addendum II to the Tautog FMP. Addendum II addressed adjustment of the compliance schedule and developed a list of issues to be reviewed and considered in a subsequent addendum or amendment. In October 2000, the Commission directed staff to initiate development of Addendum III. The primary focus of this addendum will be to provide states with additional flexibility in order to achieve the plan goals and objectives.

**II. Status of Stocks**

Tautog is a long lived species, with individuals over age 30 reported from Rhode Island and Connecticut. Most females mature (80%) at age 3. Natural mortality (M) has been estimated at  $M=0.15$  for males and  $M=0.2$  for females.

An assessment was most recently prepared in 2001, using data from 1981 through 2000. A coastwide estimate of fishing mortality rates was derived with a VPA using fisheries dependent and independent data (independent data from Massachusetts through New Jersey only). Results indicate that fishing mortality rates have declined from a high of 0.71 in 1993, to 0.41 in 2000. Since the 2000 rate exceeds both the final plan target of  $F=M=0.15$  tautog are considered overfished. Abundance indices through 2000 show a slight increase in biomass and recruitment in recent years.

For states south of NJ, a lack of data hampers efforts to estimate current fishing mortality rates and tautog abundance at a regional level of assessment. Most states are now collecting additional age and growth data with a goal of maintaining reliable F estimates. In June 2001 the Plan Review Team reviewed state compliance reports and compiled additional data that has been collected since the approval of the FMP.

**III. Status of Fishery**

The tautog fishery is primarily recreational, extending from Maine to Virginia. Most landings occur in state waters between Cape Cod and the Chesapeake Bay. Tautog have historically ranked seventh in recreational species sought in both the North Atlantic and Mid-Atlantic sub-regions. Tautog are most frequently caught in the spring and fall months, although some Mid-Atlantic Region fishermen pursue them year-round and there is an active fishery off the Virginia Coast in Winter.

Recreational landings declined from 1987 to 2000 (figure 1). Most recreationally landed tautog were caught in Massachusetts, New York, and New Jersey over the 1981 to 2000 time period, with those states accounting for 58 % of the recreational harvest by weight (table 1).

Commercial landings averaged 8% of total landings in 1999. Commercial interest in tautog has increased in response to higher market prices, exceeding \$1.00/LB at times. Most commercial landings are taken by otter trawls, with gillnets, handlines, fish pots, and lobster traps all accounting for a share.

Commercial landings fluctuated without trend around the 200,000 pound mark from 1950 until 1980. Landings began to increase in the early 1980's and reached a high of 1,157,100 pounds in 1987 (figure 2). From 1986 through 1992 landings were sustained at around a million pounds. A steep decline in landings began in 1991 and continues to the present. Commercial landings were only 208,412 pounds in 1999. Table 2 shows commercial landings from 1981 - 1999.

Recent declines in landings of both user groups can be attributed to both stock conditions and more rigorous management measures.

#### **IV. Status of Assessment Advice.**

The Tautog assessment was reviewed through an internal ASMFC stock assessment in Summer 2001. After reviewing all results, the Technical Committee recommended that fishing mortality rates be reduced by 63% to meet the final plan target ( $F=0.15$ ) and to begin rebuilding the stock.

#### **IV. Status of Research and Monitoring**

Length and abundance data are collected in trawl surveys in Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Maryland and Virginia. New York, Rhode Island, and New Jersey are collecting length and age data from party boats. Age/length information is also being collected in Massachusetts, Connecticut, Virginia, and Maryland.

Connecticut conducted a hook and release mortality study in 1993 and 1995; Virginia conducted a study in 1995. Results indicate about a 2.5% discard mortality rate. This rate was used in the last assessment.

A genetic study of Tautog from Rhode Island evaluated genetic variation patterns through DNA analysis to determine if Tautog is from one or multiple genetic stocks. This study could not determine that Tautog populations from Rhode Island to Virginia were from multiple stocks (Orbacz, Gaffney 1999).

## **V. Status of Management Measures and Issues**

The approved FMP includes commercial and recreational minimum sizes, and catch restrictions to reach the target fishing mortality. The fishing mortality target is  $F=0.15$ . Recreational fisheries will be restrained by possession limits and seasonal closures, and commercial plans will be developed by each state. The existing size limits were phased-in over a two year period.

In 1998 the PRT expressed concern about Rhode Island's recreational fishery management program. In November 1999 the ASMFC found Rhode Island out of compliance with the Interstate Fishery Management Plan for Tautog in that the state had implemented a different bag limit for its for hire fisheries and was unable to demonstrate that the alternative management program achieved the fishing mortality reduction established by the FMP. On April 4, 2000 the ASMFC was notified that Rhode Island had implemented the regulations necessary to come into compliance with the FMP. As a result, the ASMFC withdrew determination of non-compliance on April 28, 2000.

The Tautog Plan Review Team reviewed the states' annual compliance reports during summer 2001. The Plan Review Team recommended that all of the states be found in compliance.

## **VI. Current State-by-State Implementation of FMP Compliance Requirements as of June 1, 2000.**

States are expected to comply with the following provisions:

1. Submission of plans to meet  $F=0.15$  target 10/01/2001
2. Implement regulations to meet  $F=0.15$  target 4/1/2002

## **VII. Prioritized Research Needs (As of July 1, 2001)**

**\*Updated Annually (Fall of each year)**

1. Increased length sampling from the commercial/recreational fishery for all states from Massachusetts through Virginia.
2. Establish standardized state-by-state long-term fisheries independent surveys to monitor tautog abundance and length-frequency distributions, and to develop young-of-the-year indices.

3. Initiate biological sampling of the commercial catch for each gear type over the entire range of the stock (Including weight, lengths, lfd=s, age, sex, and discards).
4. Collect effort data for determining commercial and recreational CPUE.
5. Explore possible regional and local genetic differences (stock differentiation) and relate these to recruitment, growth, exploitation rates, and habitat differences. These differences can help support appropriate region-specific management strategies.
6. Continue to sample hard parts, at levels commensurate with the FMP, for annual aging from the catches of recreational and commercial fisheries and fishery-independent surveys throughout the range of the stock.
7. Define the status (condition and extent) of optimum or suitable juvenile habitats and trends in specific areas important to the species.
8. Define the susceptibility of juveniles to coastal/anthropogenic contamination and resulting effects. The synergistic effects of leaked fuel, bilge water, treated pilings, and antifouling paints on tautog health should also be studied.
9. Define the specific spawning and pre-spawning aggregating areas and wintering areas of juveniles and adults used by all major local populations, as well as the migration routes used by tautog to get to and from spawning and wintering areas and the criteria or times of use.
10. Define local and regional movement patterns and site fidelity in the southern part of the species range. This information may provide insights into questions of aggregation vs. recruitment to artificial reef locations. (Note: This work is currently being conducted as a Master=s Thesis at VIMS)
11. Reexamine the source of offshore eggs and larvae (in situ or washed out coastal spawning).
12. Confirm that tautog, like cunner, hibernate in the winter, and in what areas and temperature thresholds, for how long, and are there special habitat requirements during these times that should be protected or conserved from damage or disturbance. This information will aid in understanding behavior variability and harvest availability.
13. Conduct studies in areas where the availability of primary prey, such as blue mussels or crabs, is dependent on annual recruitment, the effect of prey recruitment variability be investigated as a factor in tautog movements (to find better prey fields), mortality (greater predation exposure when leaving shelter to forage open bottom), and relationship between reef prey availability/quality on tautog condition/fecundity.

14. Determine pot and trap escape vent dimensions needed to release tautog over a range of sizes.
15. Define larval diets and prey availability requirements. This information can be used as determinants of recruitment success and habitat function status. Information can also be used to support aquaculture ventures with this species.

Table 1. Tautog recreational harvest (A + B1) in weight (lbs.) of fish, 1981-2000 by state. Data from MRFSS as of 5/28/01.

Year	CT	DE	MD	MA	NJ	NY	RI	VA	Total
1981	242,339	6,585	10,295	790,611	161,423	1,496,039	664,568	742,653	<b>4,114,513</b>
1982	610,608	428,036	90,644	3,226,869	1,241,155	1,674,949	777,931	271,920	<b>8,322,112</b>
1983	458,581	4,438	6,550	1,837,263	414,956	1,124,844	615,592	1,267,164	<b>5,729,388</b>
1984	733,711	95,739	79,110	733,876	717,260	541,805	1,809,822	669,870	<b>5,381,193</b>
1985	471,185	144,858	1,107	328,042	741,656	2,034,903	277,385	298,796	<b>4,297,932</b>
1986	838,345	264,744	10,049	7,862,585	2,132,571	2,833,206	2,042,584	918,139	<b>16,902,223</b>
1987	1,106,606	387,075	266,093	1,751,372	2,130,955	2,288,075	507,424	442,750	<b>8,880,350</b>
1988	610,172	249,803	446,947	2,255,930	1,331,832	2,380,285	612,123	1,410,003	<b>9,297,095</b>
1989	1,038,217	743,338	78,391	1,076,365	1,289,186	1,018,016	296,889	806,337	<b>6,346,739</b>
1990	199,999	142,627	59,720	895,326	1,256,488	1,980,289	389,579	229,442	<b>5,153,470</b>
1991	648,633	354,497	106,222	798,890	2,189,144	2,352,646	1,007,548	619,215	<b>8,076,795</b>
1992	1,048,638	183,855	159,730	1,668,485	2,485,693	1,199,558	656,713	255,996	<b>7,658,668</b>
1993	531,024	217,881	105,232	752,598	1,361,612	1,800,794	389,734	758,409	<b>5,917,284</b>
1994	417,439	152,034	177,358	373,188	330,551	585,037	328,668	1,101,129	<b>3,465,404</b>
1995	402,617	793,339	115,993	309,224	1,722,714	369,643	237,094	613,348	<b>4,563,972</b>
1996	245,817	158,751	26,484	397,284	1,123,173	193,046	248,840	778,314	<b>3,171,709</b>
1997	84,297	204,419	182,995	166,042	483,639	331,530	301,109	319,257	<b>2,073,288</b>
1998	231,622	257,347	27,648	96,694	41,431	208,743	316,338	273,516	<b>1,453,339</b>
1999	61,142	358,329	37,677	363,472	511,672	761,447	223,762	203,249	<b>2,520,750</b>
2000	55,957	355,141	53,876	428,259	1,767,979	244,142	190,387	184,430	<b>33,280,171</b>
<b>Mean</b>	<b>10,036,949</b>	<b>5,502,836</b>	<b>2,042,121</b>	<b>26,112,375</b>	<b>23,435,090</b>	<b>25,418,997</b>	<b>11,894,090</b>	<b>12,163,937</b>	<b>146,606,395</b>

Table 2. Tautog commercial landings in thousands of pounds, 1981-1999 by state. Data from NMFS commercial statistics homepage as of 5/28/01.

<b>Year</b>	<b>CT</b>	<b>DE</b>	<b>MD</b>	<b>MA</b>	<b>NJ</b>	<b>NY</b>	<b>RI</b>	<b>VA</b>	<b>Total</b>
<b>1981</b>	20,500	1,000	1,200	103,000	54,400	81,400	69,800	700	<b>332,000</b>
<b>1982</b>	21,200	800	100	69,300	148,200	90,400	86,300	2,600	<b>418,900</b>
<b>1983</b>	33,500	800	-	57,600	100,600	88,400	142,600	1,700	<b>390,900</b>
<b>1984</b>	32,700	1,400	2,600	68,100	129,700	102,500	334,700	1,200	<b>672,900</b>
<b>1985</b>	49,900	3,200	2,400	63,200	125,500	84,500	403,200	1,639	<b>733,539</b>
<b>1986</b>	103,900	300	2,600	165,800	100,700	201,300	363,100	1,800	<b>939,500</b>
<b>1987</b>	159,200	500	3,800	250,000	95,200	225,200	420,500	2,700	<b>1,157,100</b>
<b>1988</b>	112,100	600	6,100	277,100	88,000	255,000	328,900	2,800	<b>1,070,600</b>
<b>1989</b>	99,706	-	3,978	352,507	51,842	285,422	214,928	7,387	<b>916,064</b>
<b>1990</b>	82,008	500	4,599	289,074	99,112	181,543	211,084	5,151	<b>873,071</b>
<b>1991</b>	54,000	1,300	3,164	354,346	93,022	226,413	371,597	5,058	<b>1,108,900</b>
<b>1992</b>	65,700	200	4,058	292,291	116,332	169,011	359,767	4,389	<b>1,011,748</b>
<b>1993</b>	44,000	300	1,432	160,336	153,474	89,467	201,593	2,660	<b>653,262</b>
<b>1994</b>	43,000	400	1,718	37,399	162,641	71,375	130,719	10,315	<b>457,567</b>
<b>1995</b>	20,466	600	4,416	35,298	116,123	72,879	95,019	27,701	<b>372,502</b>
<b>1996</b>	33,327	-	3,622	32,579	89,435	105,466	64,876	26,137	<b>322,115</b>
<b>1997</b>	14,519	841	7,663	64,275	49,726	78,228	39,601	25,471	<b>280,324</b>
<b>1998</b>	6,905	1,715	5,682	91,424	42,426	69,004	20,327	14,770	<b>252,253</b>
<b>1999</b>	12,961	844	6,489	75,685	27,392	38,033	26,107	20,901	<b>208,412</b>
<b>Mean</b>	<b>1,009,592</b>	<b>3,400</b>	<b>64,321</b>	<b>2,839,314</b>	<b>1,843,825</b>	<b>2,515,541</b>	<b>3,884,718</b>	<b>165,079</b>	<b>12,171,657</b>



Figure 1. Tautog recreational harvest (A + B1) in weight (lbs.) of fish, 1981-2000.





Figure 2. Tautog commercial landings in thousands of pounds, 1981-1999.



### 2000 Commercial Tautog Regulations

STATE	SIZE LIMIT	POSSESSION LIMITS	SEASONS	QUOTA	GEAR RESTRICTIONS
Massachusetts	16"	40	April 15-May 15 July 11-October 31		Yes
Rhode Island	16"	20	April 15-May 31	Yes	Yes
	16"	20	August 1-September 15		
	16"	20	October 15-December 31		
Connecticut	14"	a	January 1-April 30 June 15-Dec 31		Yes
New York	14"	b	some gears		Yes
New Jersey	14"		April 1-June 15 October 13-December 31	Yes	Yes
Delaware	14"/15"	See recreational	See Recreational		
Maryland	14"	5			Yes
Virginia	14"		Jan 1- April 30 Sept 1- Dec 31		Yes

### 2000 Recreational Tautog Regulations

STATE	SIZE LIMIT	POSSESSION LIMITS	SEASONS
Massachusetts	16"	6	-
Rhode Island	16"	3	May 1-October 14
	16"	10	October 15-December 31
Connecticut	14"	4 (daily)	Jan 1-Apr 30, June 15- Dec 31
New York	14"	c	-
New Jersey	14"	d	d
Delaware	14"	10 <sup>e</sup>	Jul 1 - Mar 31
	15"	3	Apr 1 - Jun 30
Maryland	14"	5	-
Virginia	14"	7	Open

- a The trawl fishery has a possession limit of 75 fish, the commercial hook, fish pot, trap net, fyke net, and gill net fisheries the possession limit is 25 fish, and in the lobster pot fishery the possession limit is 10 fish. Holders of Connecticut Marine Pound Net Registration may possess up to twelve fish year round except that during the May 1 through June 14 closed season all female tautog must be released without avoidable injury. All possession limits are daily limits.
- b New York has a 25 fish vessel trip limit for commercially caught tautog, except only 10 per vessel are allowed when lobster pot gear and more than six lobsters are in possession.
- c New York has a one fish bag limit in effect for recreational catches from June 1 through October 6 and a 10 fish bag limit in effect from October 7 through May 31.
- d New Jersey has a possession limit of 10 tautog from October 10 through May 31. The possession limit from June 1 through October 9 is 1 tautog.
- e Delaware has an 11 day closure from September 8 through September 18.