


2013 REVIEW OF THE
ATLANTIC STATES MARINE FISHERIES COMMISSION
FISHERY MANAGEMENT PLAN FOR

ATLANTIC CROAKER
(Micropogonias undulatus)

2012 FISHING YEAR



Atlantic Croaker Plan Review Team

Joe Grist, Virginia Marine Resources Commission
Wilson Laney, Ph.D., United States Fish and Wildlife Service
Chris McDonough, South Carolina Department of Natural Resources
Jason Rock, North Carolina Department of Marine Fisheries
Kirby Rootes-Murdy, Atlantic States Marine Fisheries Commission, Chair

Table of Contents

| | | |
|-------|---|----|
| I. | Status of the Fishery Management Plan..... | 1 |
| II. | Status of the Stock | 2 |
| III. | Status of the Fishery..... | 3 |
| IV. | Status of Assessment Advice | 4 |
| V. | Status of Research and Monitoring..... | 5 |
| VI. | Status of Management Measures and Issues..... | 6 |
| VII. | Implementation of FMP Compliance Requirements for 2009..... | 8 |
| VIII. | Recommendations..... | 8 |
| IX. | References..... | 10 |
| X. | Figures..... | 11 |
| XI. | Tables..... | 13 |

I. Status of the Fishery Management Plan

| | |
|----------------------------------|--|
| <u>Date of FMP Approval:</u> | Original FMP – October 1987 |
| <u>Amendments:</u> | Amendment 1 – November 2005 (implemented January 2006) Addendum I – March 2011 |
| <u>Management Areas:</u> | The Atlantic coast distribution of the resource from New Jersey through Florida |
| <u>Active Boards/Committees:</u> | South Atlantic State/Federal Fisheries Management Board; Atlantic Croaker Technical Committee, Stock Assessment Subcommittee, and Plan Review Team; South Atlantic Species Advisory Panel |

The Fishery Management Plan (FMP) for Atlantic Croaker was adopted in 1987 and included the states from Maryland through Florida (ASMFC 1987). Subsequently, the South Atlantic State/Federal Fisheries Management Board (Board) reviewed the FMP and found its recommendations to be vague and recommended that an amendment be prepared to define management measures necessary to achieve the goals of the FMP. The Interstate Fisheries Management Program Policy Board also adopted the finding that the original FMP did not contain any management measures that states were required to implement.

In 2002, the Board directed the Atlantic Croaker Technical Committee to conduct the first coastwide stock assessment of the species in preparation of developing an amendment. The Atlantic Croaker Stock Assessment Subcommittee developed a stock assessment in 2003, which was approved by a Southeast Data Assessment Review (SEDAR) panel for use in management in June 2004 (ASMFC 2005a). The Board quickly initiated the development of an amendment. In November 2005, the Board approved Amendment 1 to the Atlantic Croaker FMP (ASMFC 2005b). The amendment was fully implemented by January 1, 2006.

The goal of Amendment 1 is to utilize interstate management to perpetuate the self-sustainable Atlantic croaker resource throughout its range and generate the greatest economic and social benefits from its commercial and recreational harvest and utilization over time. Amendment 1 contains four objectives:

- 1) Manage the fishing mortality rate for Atlantic croaker to provide adequate spawning potential to sustain long-term abundance of the Atlantic croaker population.
- 2) Manage the Atlantic croaker stock to maintain the spawning stock biomass above the target biomass levels and restrict fishing mortality to rates below the threshold.
- 3) Develop a management program for restoring and maintaining essential Atlantic croaker habitat.
- 4) Develop research priorities that will further refine the Atlantic croaker management program to maximize the biological, social, and economic benefits derived from the Atlantic croaker population.

Amendment 1 expanded the management area to include the states from New Jersey through Florida. Consistent with the stock assessment completed in 2004, the amendment defined two Atlantic coast management regions: the south-Atlantic region, including the states Florida

through South Carolina; and the mid-Atlantic region, including the states North Carolina through New Jersey.

Amendment 1 established biological reference points (BRPs) to define overfished and overfishing stock status for the mid-Atlantic region only. Reliable stock estimates and BRPs for the South Atlantic region could not be developed during the 2004 stock assessment due to a lack of data. The BRPs were based on maximum sustainable yield (MSY), and included threshold and target levels of fishing mortality (F) and spawning stock biomass (SSB): F threshold = F_{MSY} (estimated to be 0.39); F target = $0.75 \times F_{MSY}$ (estimated to be 0.29); SSB threshold = $0.7 \times SSB_{MSY}$ (estimated to be 44.65 million pounds); and SSB target = SSB_{MSY} (estimated to be 63.78 million pounds). An SSB estimate below the SSB threshold results in an overfished status determination, and an F estimate above the F threshold results in an overfishing status determination. The Amendment established that the Board would take action, including a stock rebuilding schedule if necessary, should the BRPs indicate an overfished stock or a stock subject to overfishing.

Amendment 1 did not require any specific measures restricting recreational or commercial harvest of Atlantic croaker. States with more conservative measures were encouraged to maintain those regulations (Table 1). Through adaptive management, the Management Board may revise Amendment 1, and regulatory and/or monitoring requirements could be included in the resulting addendum, along with procedures for determining de minimis status and implementing alternative management programs via conservation equivalency.

The Board initiated Addendum I to Amendment I at its August 2010 meeting, following the updated stock assessment, in order to address the proposed reference points and management unit. The stock assessment evaluated the stock based on a coastwide unit, rather than the two management units established within Amendment I. In approving the final Addendum I, the Management Board approved the consolidation of the stock into one management unit, as proposed by the stock assessment. In addition, Addendum I established a procedure, similar to other species, by which the Board may approve peer-reviewed BRPs without a full administrative process, such as an amendment or addendum.

Addendum I did not add or change any additional management measures or requirements. The only existing requirement is for states to submit an annual compliance report by July 1 of each year that contains commercial and recreational landings as well as results from any monitoring programs that intercept Atlantic croaker.

II. Status of the Stock

Stock status is based on the data and results of the 2010 stock assessment (ASMFC 2010). Results include revised biological reference points (below). These reference points are ratio-based and apply to the entire coastwide resource (unlike those in Amendment 1). Overfishing is occurring if F/F_{MSY} is greater than 1 and the stock is considered overfished if $SSB/(SSB_{MSY}(1-M))$ is less than 1.

| | Overfishing Definition | Overfished Definition |
|-----------|--------------------------|----------------------------|
| Target | $F/(F_{MSY} * 0.75) = 1$ | $SSB/SSB_{MSY} = 1$ |
| Threshold | $F/F_{MSY} = 1$ | $SSB/(SSB_{MSY}(1-M)) = 1$ |

Atlantic croaker is not experiencing overfishing. Biomass has been increasing and fishing mortality decreasing since the late 1980s. Biomass conclusions are based on information from the data compiled for the assessment, namely increasing indices of relative abundance and expanding age structure in the catch and indices. Model estimated values of fishing mortality (F), spawning stock biomass (SSB), and biological reference points are too uncertain to be used to determine stock status. However, the ratio of F to F_{MSY} (the F needed to produce maximum sustainable yield) is reliable and can be used to determine that overfishing is not occurring. It is not possible to be confident with regard to stock status, particularly a biomass determination, until the discards of Atlantic croaker from the South Atlantic shrimp trawl fishery can be adequately estimated and incorporated into the stock assessment.

Absolute estimates of total F are unavailable because of model uncertainty; however, the general trend in total F from the model is considered reliable due to support from the data. The trend in total F decreases substantially during the first five years of the time series (1988-1992) and shows an overall decline over the remainder of the time series, except for occasional, brief spikes (Figure 1). Retrospective analysis of the model showed that estimates of F decreased as more years of data were used. A series of sensitivity runs conducted over a range of plausible values of shrimp-trawl fishing mortality found that the ratio of directed fishing mortality to F_{MSY} was less than one in all cases, indicating overfishing was not occurring.

Absolute estimates of SSB are unavailable because of model uncertainty; however, the general trend in SSB from the model is considered reliable due to support from the data. Spawning stock biomass shows a nearly consistent increasing trend since 1998 (Figure 2). Sensitivity runs of the model, including rough estimates of shrimp trawl discards, do not change the overall trend in SSB. Retrospective analysis of the model showed that estimates of SSB increased as more years of data were used.

Recruitment, estimated in the model as age-1 abundance, has been variable but generally increasing over the time series. Figure 2 shows the trend in recruitment; absolute values are omitted because of uncertainty in abundance estimates. The model estimated the production of strong year classes in 1997, 2001, and 2007.

III. Status of the Fishery

Total Atlantic croaker harvest from New Jersey through the east coast of Florida in 2012 is estimated at 14.6 million pounds (Tables 2 and 3, Figure 3). This represents a 35 percent decline in total harvest since the peak at 41.2 million pounds in 2001 (61% commercial decline, 73% recreational decline). The commercial and recreational fisheries harvested 80 and 20 percent of the total, respectively. The vast majority of landings are from the Mid-Atlantic region (97% in 2012), and the recent decline in total landings is a result of both commercial and recreational landings declines in that region, although some states showed increases in either or both sectors (Figure 4). Commercial and recreational landings in the South Atlantic region have been generally stable over the last decade; however, 2010 showed large decreases in the recreational

harvest of the South Atlantic states' fisheries, though nothing of the same magnitude as in the Mid-Atlantic states. Recreational and commercial harvests in the South Atlantic region rebounded to previous levels in 2011 and held stable in 2012.

Atlantic coast commercial landings of Atlantic croaker exhibit a cyclical pattern, with low domains in the 1960s to early 1970s and the 1980s to early 1990s, and high domains in the mid-to-late 1970s and the mid-1990s to the present (Figure 3). This cyclical pattern was noted in the recent 2010 stock assessment, noting that the 50-year time series follows this pattern and that the current trend has been towards a low. Commercial landings increased from a low of 3.7 million pounds in 1991 to 30.1 million pounds in 2001 (Table 2); however, landings have declined consistently since 2003 to 11.6 million pounds in 2012, which registers below the 1960-2011 average of 13.6 million pounds. Within the management unit, the majority of 2012 commercial landings came from Virginia (59%) and North Carolina (27%). Maryland had the next highest level, with 8% of the coastwide landings.

From 1981-2012, recreational landings of Atlantic croaker from New Jersey through Florida have varied between 2.8 million fish (1.3 million pounds) and 13.2 million fish (11.1 million pounds; Tables 3 and 4, Figure 5). Landings general increased until 2001, held stable from 2001-2006 before exhibiting a declining trend from 2007 through 2012. The 2012 landings are estimated at 5.4 million fish and 2.9 million pounds, continuing the decline from 2009. Virginia was responsible for 65% of the 2012 recreational landings, in numbers of fish, followed by Maryland (13%), and Florida (11%). The number of recreational releases has increased over the time series, with a short decline from 2009-2011 (Figure 5). In 2012, anglers released 10.5 million fish, which is less than the ten-year (2002-2011) average of 11.9 million fish (Table 5). Anglers released an estimated 66% of the croaker catch in 2012 (Figure 5).

IV. Status of Assessment Advice

A statistical catch-at-age (SCA) model was used in the last Atlantic croaker stock assessment (ASMFC 2010). This model combines the catch-at-age data from the commercial and recreational fisheries with information from fishery-independent surveys and biological information such as growth rates and natural mortality rates to estimate the size of each age class and the exploitation rate of the population. The assessment was peer reviewed by a panel of experts in conjunction with the Southeast Data, Assessment, and Review (SEDAR) process.

The Review Panel was unable to support some of the assessment results due to uncertainty regarding the estimation of Atlantic croaker discards in the shrimp trawl fishery, and the application of estimates in modeling. Specifically, model-estimated values of stock size, fishing mortality, and biological reference points are too uncertain for use; however, the trends in model-estimated parameters and ratio-based fishing F reference points are considered reliable. Adequate discard estimates cannot be developed from currently available data, and assessments of Atlantic croaker will be unreliable until adequate estimates are properly incorporated into modeling. Despite the uncertainty in assessment results caused by shrimp trawl bycatch, the Review Panel concluded that it is unlikely that the stock is in trouble. The stock is not experiencing overfishing, biomass has been trending up, commercial catches are stable, and discards from the shrimp trawl fishery have been much reduced.

V. Status of Research and Monitoring

There are no research or monitoring programs required of the states except for the submission of an annual compliance report. The following fishery-dependent (other than catch and effort data) and fishery-independent monitoring programs were reported in the 2013 compliance reports.

Fishery-Dependent Monitoring

- New Jersey: commercial fishery biological sampling since 2006 (660 length measurements and 619 otolith ages in 2012)
- Maryland: commercial pound net fishery biological sampling (1,842 length measurements, 255 otoliths collected in 2012), at-sea sampling supplemented with dealer sampling beginning in 2009 (571 length and weight measurements in 2012); Maryland Charter Boat CPUE (1993-present; 2012 CPUE continued decline from 2010 but still above time-series mean)
- Virginia: commercial fishery biological sampling (7,121 length and weight measurements, 400 otolith ages, and 672 sex determinations in 2012)
- North Carolina: commercial fishery biological sampling since 1982 for length (2011 n = 7,098), weight (2011 n = 14,829), otolith, sex determination, and reproductive condition
- South Carolina: recreational fishery biological sampling via state finfish survey (137 length measurements in 2012)
- Georgia: recreational fishery biological sampling via carcass collections (1 fish in 2012)
- Florida: commercial fishery biological sampling (43 length measurements in 2012)

Fishery-Independent Monitoring

- New Jersey: nearshore ocean (within 12 nm) juvenile trawl surveys (1988-present; 2012 CPUE well above time-series average; nearshore Delaware Bay juvenile trawl survey (1991-present; 2012 CPUE low but near time-series average); Delaware River juvenile seine survey (1980-present; 2012 CPUE high and above time-series average)
- Delaware: offshore Delaware Bay adult finfish trawl survey (1966-present; 2012 n = 8,885; 662% increase in catch per nm towed over 2011 but still below time-series mean); nearshore Delaware Bay and River juvenile finfish trawl survey (1980-present; 2012 index (geometric mean) declined 4% from 2011 and fell below time-series mean)
- Maryland: Atlantic coast bays juvenile otter trawl survey (standardized from 1989-present); Chesapeake Bay juvenile trawl survey (standardized from 1989-present; 2012 CPUE increased from 2011, recording the seventh highest value in 24 year time series); incidental catches in Maryland coastal bays juvenile seine survey (1972-present) and Chesapeake Bay juvenile seine survey (1959-present; 2012 indices increased or were stable relative to 2011)
- Virginia: VIMS Juvenile Finfish and Blue Crab Trawl Survey (1988-present; 2012 index representing the 2011 year class, is the third highest on record, from 1988 to 2012, for both the 'mean all' and 'mean rivers' indices.)
- North Carolina: Pamlico Sound juvenile trawl survey (1987-present; 2012 juvenile abundance index was the second highest recorded in NC and was well above time-series mean)
- South Carolina: estuarine electroshock survey for juveniles (1991-present; 2012 CPUE decreased 50% , marking the fourth year in a row of being below the long term mean); SEAMAP shallow water (15-30 ft) trawl survey from Cape Hatteras to Cape Canaveral (1989-present; 2012 CPUE decreased 31.1% but remained above time-series mean); inshore estuarine trammel net survey for adults (May-September, 1991-present; 2012

CPUE increased 53% from 2011 and catch effort in 2012 was just below the long term mean)

- Georgia: Marine Sportfish Population Health Survey (trammel and gill net, 2002-present; 2012 n = 158); Ecological Monitoring Survey (trawl, 2003-present; 2012 n = 7,508; CPUE declined by 50% from 2011)
- Florida: juvenile seine survey (1996-present; 2012 index continued variable trend with a decrease from 2011); juvenile trawl survey (2002-present; 2012 index continued variable trend with a decrease from 2011); adult haul seine survey (2001-present; 2012 index value decreased from 2011)

The Northeast Fishery Science Center's groundfish trawl survey also samples croaker from New Jersey to Cape Hatteras. Researchers from various agencies and institutions have conducted numerous studies on Atlantic croaker. Research topics include, but are not limited to: environmental effects on recruitment, population modeling, genetic stock identification, geographic variation in life history/populations dynamics, scale-otolith age comparisons, habitat preference, and bycatch reduction gear research.

Ageing Workshop

An Atlantic Croaker Ageing Workshop was held in October 2008. Conducting a workshop to standardize the otolith sectioning and ageing procedures and the current age dataset had been a longstanding research need for Atlantic croaker, especially prior to the 2010 benchmark assessment. Representatives from New Jersey, Maryland, Virginia, North Carolina, South Carolina, Georgia and the Gulf Council attended the workshop. The resulting standardized ageing procedure was published in an ASMFC reference document, with some states having already incorporated ageing instructions into their references.

VI. Status of Management Measures and Issues

Fishery Management Plan

Amendment 1 was fully implemented by January 1, 2006, and provided the management plan for the 2009 fishing year. There are no interstate regulatory requirements for Atlantic croaker. Should regulatory requirements be implemented in the future, all state programs must include law enforcement capabilities adequate for successfully implementing the regulations. Addendum I to Amendment 1 was initiated in August 2010 and approved in March 2011, in order to 1) revise the biological reference points to be ratio-based, and 2) remove the distinction of two regions within the management unit, based on the results of the 2010 stock assessment.

De Minimis Requests

States are permitted to request *de minimis* status if, for the preceding three years for which data are available, their average commercial landings or recreational landings (by weight) constitute less than 1% of the coastwide commercial or recreational landings for the same three year period. A state may qualify for *de minimis* in either its recreational or commercial sector, or both, but will only qualify for exemptions in the sector(s) that they qualify for as *de minimis*. Amendment 1 does not include any compliance requirements other than annual state reporting, which is still required of *de minimis* states, thus *de minimis* status does not exempt states from any measures.

In the annual compliance reports, the following states requested *de minimis* status: Delaware (commercial fishery), South Carolina (commercial fishery), Georgia (commercial and recreational fisheries), and Florida (commercial fishery). The commercial and recreational *de minimis* criteria for 2011 are based on 1% of the average coastwide 2010-2012 landings in each fishery: 129,039 pounds for the commercial fishery and 35,135 pounds for the recreational fishery. The Delaware commercial fishery qualifies for *de minimis* status with an average of 6,727 pounds. The South Carolina commercial fishery qualifies for *de minimis* status with an average of 36 pounds. The Georgia commercial and recreational fisheries qualify for *de minimis* status with averages of less than 1,000 pounds (confidential) and 15,039 pounds, respectively. The Florida commercial fishery qualifies for *de minimis* status with an average of 50,386 pounds.

Bycatch Reduction

Atlantic croaker is subject to both direct and indirect fishing mortality. Historically, croaker ranked as one of the most abundant species in the bycatch of the south Atlantic shrimp trawl fishery. As a result, the original FMP recommended that bycatch reduction devices (BRDs) be developed and required in the shrimp trawl fishery. Since then the states of North Carolina through Florida have all enacted requirements for the use of BRDs in shrimp trawl nets in state waters, and croaker bycatch from this fishery has been reduced (ASMFC 2010). However, monitoring of bycatch and discards from this fishery is inadequate and results in the major source of uncertainty for assessing this stock, as well as other important Mid- and South Atlantic species. Most of the discarded croakers are age-0 and thus likely have not yet reached maturity (ASMFC 2010). North Carolina Department of Marine Fisheries has secured funding for a two-year study, beginning in 2012, to collect bycatch data from state shrimp trawlers. These data will be valuable for incorporating estimates of removals in the next stock assessment.

Atlantic croaker are also discarded from other commercial fishing gears. This is primarily due to market pressures and few restrictions on croaker harvest at the state level. The NMFS Pelagic Observer Program provides data to estimate these discards for use in assessments; however, the time series is limited and only discards from gill nets and otter trawls could be estimated for the last assessment based on the available data. Since 1988, estimated discards have fluctuated between 94 and 15,176 mt without trend, averaging 2,503 mt (ASMFC 2010).

Atlantic croaker has also been a major component of the scrap/bait fishery. Landings from this fishery are not reported to the species level, except for North Carolina, which has a continuous program in place to sample the landings and enables estimating scrap landings of croaker for use in the stock assessment. As part of the recent stock assessment, North Carolina estimated the scrap/bait landings, which have declined in recent years, from a high of 1,569 mt in 1989 to a low of 84 mt in 2008, primarily due to restrictions placed on the fisheries that produced the highest scrap/bait landings (ASMFC 2010). Several of the regulations instituted by North Carolina include a ban on flynet fishing south of Cape Hatteras, incidental finfish limits for shrimp and crab trawls in inside waters, minimum mesh size restrictions in trawls, and culling panels in long haul seines. Monitoring programs are needed to account for bait/scrap landings in other states.

Several states have implemented other commercial gear requirements that further reduce bycatch and bycatch mortality, while others continue to encourage the use of these BRD devices. NOAA

Fisheries recently published a notice on June 24, 2011 for public scoping in the Federal Register to expand the methods for reducing bycatch interactions with sea turtles, which may have additional effects on the bycatch of finfish like Atlantic croaker in trawls (76 FR 37050). Continuing to reduce the quantity of sub-adult croaker harvested should increase spawning stock biomass and yield per recruit.

Atlantic croaker are also subject to recreational discarding. The number of Atlantic croaker released alive by recreational anglers has generally increased over time. Ten percent of croakers released alive were estimated to die as a result of being discarded for the last stock assessment (ASMFC 2010). The use of circle hooks and appropriate handling techniques can help to reduce mortality of released fish.

Trigger Exercises

Amendment 1 requires the Technical Committee to conduct stock assessments every five years unless prompted by the annual trigger exercise. The primary hard trigger is based on landings data; however, catch-per-unit-effort (CPUE) will become the premier trigger when the quality and quantity of these data improve. A stock assessment will be triggered if the most recent year's commercial or recreational landings are less than 70% of the previous two years' average landings (ASMFC 2005b).

In 2011, the recreational landings dropped to 75.9% of the previous two-year average, therefore not triggering a stock assessment update or benchmark. The Atlantic Croaker Technical Committee reviewed the triggers, as well as discussed development of new triggers as tasked by the Board via three conference calls in May, June, and July. While the commercial and recreational landings, along with the estimates of landings per unit effort, have shown decreases, the fishery-independent indices have not indicated major issues with the stock. Based on the available data and benefits and disadvantages of performing an update to the stock assessment, the Technical Committee recommended the Board not perform an assessment but rather allow the Technical Committee to build other approaches into the trigger exercises, such as using a traffic light methodology. The 2013 Atlantic Croaker Trigger Report further details the Technical Committee's this approach and subsequent recommendations.

VII. Implementation of FMP Compliance Requirements for 2012

The PRT finds that all states have fulfilled the requirements of Amendment 1.

VIII. Recommendations

Management and Regulatory Recommendations

- Encourage the use of circle hooks to minimize recreational discard mortality.
- Consider approval of the *de minimis* requests from Delaware, South Carolina, Georgia, and Florida.
- Consider the basic research and monitoring information needed for informed management in light of the budgetary constraints limiting all state governments
- Support the Technical Committee's recommendation to develop new assessment/management triggers for use in management by the Board

Research and Monitoring Recommendations

High Priority

- Develop and implement compatible and coordinated sampling programs for the South Atlantic shrimp trawl fishery in order to monitor and characterize Atlantic croaker bycatch in this fishery.
- Continue fisheries-independent surveys throughout the species range, with increased focus on collecting subsamples in the southern range
- Encourage fishery-dependent biological sampling, with increased focus in the southern range and expanding the commercial and recreational fishery samples to afford a full age-length key
- Determine migratory patterns and mixing rates through cooperative, multi-jurisdictional tagging studies; further studies on relative degree of genetic separation between fish in the northern and southern range of species; and continue research and analysis of otolith microchemistry data.
- Collect bio-profile information and conduct studies on growth rates, age structure, estimates of fecundity, and maturity schedule throughout the species range with a standardized protocol.
- Evaluate bycatch and discard estimates from commercial and recreational fisheries, and extend coverage of scrap fishery sampling to other states.
- Develop fishery-independent size, age, and sex specific relative abundance estimates to monitor long-term changes in croaker abundance.
- Maintain funding for current surveys and monitoring to provide needed information for stock monitoring and assessment

Medium Priority

- Develop age-size data that are representative of all seasons and areas in the fisheries on an annual basis.
- Improve catch and effort statistics from the commercial and recreational fisheries and develop more rigorous methods to standardize catch-per-unit-effort.
- Collect data on fishing attributes necessary to develop gear-type-specific fishing effort estimates.
- Evaluate commercial and recreational mortality under varying environmental factors and fishery practices and include in updated assessment.
- Update studies on the effectiveness of bycatch reduction devices (BRDs) in reducing croaker bycatch.
- Validate otolith aging methods with appropriate methods, e.g., tagging, chemical marking.
- Evaluate the optimum utilization (economic and biological) of a long-term fluctuating population such as croaker.
- Identify essential habitat requirements.
- Determine species interactions and predator/prey relationships for croaker (prey) and other more highly valued fisheries (predators).
- Determine the impacts of any dredging activity (i.e. for beach re-nourishment) on all life history stages of croaker.
- Investigate environmental covariates in stock assessment models.
- Examine socio-economic aspects of the fishery.
- Recover historical data in order to have landings data from NOAA at a finer scale

- Re-examine historical ichthyoplankton studies of the Chesapeake Bay for an indication of the magnitude of estuarine spawning.

IX. References

- Atlantic States Marine Fisheries Commission (ASMFC). 1987. Fishery Management Plan for Atlantic Croaker. Washington (DC): ASMFC. Fishery Management Report No. 10. 90 p.
- ASMFC. 2005a. Atlantic Croaker Stock Assessment & Peer Review Reports. Washington (DC): ASMFC. 370 p.
- ASMFC. 2005b. Amendment 1 to the Interstate Fishery Management Plan for Atlantic Croaker. Washington (DC): ASMFC. Fishery Management Report No. 44. 92 p.
- ASMFC. 2010. Atlantic Croaker 2010 Benchmark Stock Assessment. Washington (DC): ASMFC. 366 p.

X. Figures




Figure 1. Trend in estimated total fishing mortality rate (F) of Atlantic croaker (Absolute estimates of F are unreliable because of uncertainty regarding the estimation of Atlantic croaker discards in the shrimp trawl fishery, and the application of estimates in modeling. Source: ASMFC 2010.)




Figure 2. Trends in estimated spawning stock biomass (SSB, metric tons) and age-1 recruitment (numbers of fish) of Atlantic croaker (Absolute estimates of stock size are unreliable because of uncertainty regarding the estimation of Atlantic croaker discards in the shrimp trawl fishery, and the application of estimates in modeling. Source: ASMFC 2010.)




Figure 3. Atlantic croaker commercial, recreational, and total landings (pounds)
 (See Tables 2 and 3 for values and source information. Commercial landings estimate for 2012 is preliminary. Reliable recreational landings estimates are not available before 1981.)




Figure 4. Mid-Atlantic (NJ-NC) and South Atlantic (SC-FL) landings (pounds)
 (See Tables 2 and 3 for values and source information.)




Figure 5. Recreational catch (landings and alive releases, in numbers) and the percent of catch that is released, 1981-2012

(See Tables 4 and 5 for values and source information.)

XI. Tables

Table 1. Summary of state regulations for Atlantic croaker in 2012*

| State | Recreational | Commercial |
|-------|---|---|
| NJ | none | otter/beam trawl mesh restriction for directed croaker harvest (>100 lbs in possession) |
| DE | 8" minimum; recreational gill nets (up to 200 ft.) with license | 8" minimum |
| MD | 9" min, 25 fish/day, charter boat logbooks | 9" minimum; open 3/16 to 12/31 |
| PRFC | 25 fish/day | pound net season: 2/25 to 12/15 |
| VA | none | none |
| NC | recreational use of commercial gears with license and gear restrictions | |
| SC | mandatory for-hire logbooks | |
| GA | 8" min, 25 fish/day | 8" minimum; 25 fish/day limit except for shrimp trawls (no limit) |
| FL | none | none |

* A commercial fishing license is required to sell croaker in all states with fisheries. For all states, general gear restrictions affect commercial croaker harvest.

Table 2. Commercial harvest (pounds) of Atlantic croaker by state, 1981-2012

(Estimates for 2012 are preliminary. Sources: state compliance reports; personal communication with ACCSP, Arlington, VA.)

| Year | NJ | DE | MD | PRFC | VA | NC | SC | GA | FL | Total |
|------|-----------|--------|-----------|-----------|------------|------------|-------|-------|---------|------------|
| 1981 | 23,500 | 0 | 2,104 | 648 | 429,800 | 11,205,342 | 2,441 | 1,038 | 72,112 | 11,736,985 |
| 1982 | 100 | 0 | 7,091 | 188 | 119,300 | 10,824,953 | 386 | 2,177 | 95,357 | 11,049,552 |
| 1983 | 200 | 0 | 417 | 1,549 | 150,400 | 7,249,680 | 3,200 | 1,097 | 81,737 | 7,488,280 |
| 1984 | 57,700 | 0 | 27,072 | 73,701 | 817,700 | 9,170,775 | 3,793 | 434 | 131,375 | 10,282,550 |
| 1985 | 48,800 | 100 | 9,510 | 19,854 | 2,171,821 | 8,714,432 | 1,256 | | 153,803 | 11,119,576 |
| 1986 | 106,000 | 500 | 135,922 | 99,373 | 2,367,000 | 9,424,828 | 924 | | 173,531 | 12,308,078 |
| 1987 | 357,600 | 800 | 119,409 | 102,691 | 2,719,500 | 7,289,191 | 698 | 553 | 217,932 | 10,808,374 |
| 1988 | 30,100 | 200 | 98,855 | 12,796 | 1,749,200 | 8,434,415 | 2,614 | 304 | 140,033 | 10,468,517 |
| 1989 | 137,100 | 0 | 89,173 | 5,579 | 949,649 | 6,824,088 | 1,950 | | 95,021 | 8,102,560 |
| 1990 | 644 | 42 | 2,473 | 5,115 | 201,353 | 5,769,512 | 1,190 | | 104,402 | 6,084,731 |
| 1991 | 31,292 | 700 | 6,183 | 996 | 164,126 | 3,436,960 | * | | 56,739 | 3,696,996 |
| 1992 | 51,600 | 800 | 17,050 | 17,692 | 1,339,353 | 2,796,612 | | | 79,040 | 4,302,147 |
| 1993 | 183,414 | 2,500 | 114,159 | 262,482 | 5,326,293 | 3,267,652 | * | | 52,031 | 9,208,531 |
| 1994 | 117,256 | 3,000 | 158,918 | 240,271 | 5,759,975 | 4,615,754 | * | | 96,018 | 10,991,192 |
| 1995 | 334,654 | 13,000 | 489,506 | 606,184 | 6,949,639 | 6,021,284 | * | | 22,879 | 14,437,146 |
| 1996 | 621,889 | 9,681 | 792,326 | 1,427,285 | 9,409,904 | 9,961,834 | | | 26,045 | 22,248,964 |
| 1997 | 1,994,446 | 10,509 | 1,088,969 | 1,518,196 | 12,832,221 | 10,711,667 | * | | 36,577 | 28,192,585 |
| 1998 | 1,029,332 | 10,368 | 1,006,529 | 610,885 | 11,898,586 | 10,865,897 | | | 26,418 | 25,448,015 |
| 1999 | 2,071,046 | 14,729 | 948,191 | 1,190,138 | 12,481,326 | 10,185,507 | | | 26,824 | 26,917,761 |
| 2000 | 2,130,465 | 11,121 | 902,379 | 1,812,130 | 12,822,400 | 10,122,627 | | | 37,953 | 27,839,075 |
| 2001 | 1,389,837 | 22,736 | 1,488,815 | 1,963,294 | 13,214,731 | 12,017,424 | | * | 14,831 | 30,111,668 |
| 2002 | 1,828,484 | 10,732 | 894,879 | 1,421,094 | 12,133,834 | 10,189,153 | * | * | 17,191 | 26,495,367 |
| 2003 | 1,575,738 | 16,561 | 713,205 | 1,128,003 | 10,937,167 | 14,429,197 | 140 | * | 16,402 | 28,816,413 |
| 2004 | 2,067,992 | 32,729 | 1,354,982 | 1,631,596 | 8,550,574 | 11,993,003 | * | * | 11,413 | 25,642,289 |
| 2005 | 1,847,753 | 39,931 | 972,800 | 481,912 | 8,248,441 | 11,903,292 | 41 | * | 16,520 | 23,510,690 |
| 2006 | 1,617,144 | 19,277 | 466,833 | 670,276 | 9,293,410 | 10,396,554 | 160 | * | 30,272 | 22,493,926 |
| 2007 | 1,358,000 | 13,651 | 474,388 | 188,567 | 10,697,251 | 7,301,295 | * | | 27,028 | 20,060,180 |
| 2008 | 946,062 | 10,465 | 592,211 | 337,062 | 11,925,676 | 5,791,874 | 116 | * | 31,560 | 19,635,026 |
| 2009 | 585,552 | 16,258 | 433,238 | 234,101 | 8,422,147 | 6,135,427 | 75 | 0 | 32,310 | 15,859,108 |
| 2010 | 342,116 | 6,024 | 490,067 | 163,371 | 6,574,894 | 7,312,159 | 3 | 0 | 36,882 | 14,925,516 |
| 2011 | 465,049 | 11,346 | 694,673 | 238,050 | 5,379,417 | 5,054,186 | 44 | * | 44,899 | 11,933,656 |
| 2012 | 363,381 | 2,811 | 901,455 | 273,849 | 6,908,462 | 3,106,616 | 62 | * | 69,378 | 11,626,014 |

* confidential data

Table 3. Recreational harvest (pounds) of Atlantic croaker by state, 1981-2012

(Source: personal communication with NMFS Fisheries Statistics Division, Silver Spring, MD.)

| Year | NJ | DE | MD | VA | NC | SC | GA | FL | Total |
|------|-----------|---------|-----------|-----------|---------|---------|---------|-----------|------------|
| 1981 | 582 | 2,317 | | 535,297 | 426,240 | 67,284 | 9,665 | 305,547 | 1,346,932 |
| 1982 | | | 70,276 | 455,250 | 264,607 | 67,015 | 45,161 | 754,956 | 1,657,265 |
| 1983 | | | 32,053 | 486,006 | 395,402 | 14,158 | 25,412 | 510,599 | 1,463,630 |
| 1984 | | | 86,462 | 634,870 | 584,660 | 161,661 | 80,684 | 1,856,599 | 3,404,936 |
| 1985 | | | 17,169 | 843,414 | 278,214 | 72,780 | 40,421 | 684,449 | 1,936,447 |
| 1986 | | 2,595 | 116,542 | 2,034,337 | 126,888 | 173,028 | 21,504 | 2,783,651 | 5,258,545 |
| 1987 | | | 191,628 | 1,306,814 | 352,346 | 64,696 | 14,947 | 1,005,053 | 2,935,484 |
| 1988 | | 827 | 926,399 | 2,390,573 | 935,460 | 54,313 | 20,313 | 316,900 | 4,644,785 |
| 1989 | | 284 | 19,189 | 1,329,680 | 658,567 | 80,580 | 21,138 | 268,335 | 2,377,773 |
| 1990 | | 112 | 37,873 | 875,427 | 347,183 | 123,795 | 205,352 | 127,525 | 1,717,267 |
| 1991 | 4,264 | 10,972 | 117,210 | 1,728,021 | 157,660 | 16,173 | 54,116 | 460,453 | 2,548,869 |
| 1992 | | 3,291 | 53,556 | 1,768,962 | 233,533 | 28,512 | 132,596 | 407,672 | 2,628,122 |
| 1993 | 844 | 9,641 | 476,866 | 1,993,915 | 282,910 | 18,005 | 55,604 | 180,517 | 3,018,302 |
| 1994 | 818 | 2,892 | 991,166 | 3,024,118 | 351,230 | 128,306 | 34,048 | 337,474 | 4,870,052 |
| 1995 | 9,515 | 82,864 | 567,149 | 2,675,381 | 326,135 | 25,386 | 20,862 | 301,918 | 4,009,210 |
| 1996 | 39,099 | 205,526 | 702,037 | 2,716,759 | 346,501 | 14,480 | 21,797 | 50,038 | 4,096,237 |
| 1997 | 278,758 | 340,198 | 1,117,999 | 5,522,195 | 309,457 | 53,863 | 26,272 | 113,096 | 7,761,838 |
| 1998 | 135,733 | 293,560 | 1,150,459 | 5,920,436 | 161,117 | 76,821 | 30,966 | 141,756 | 7,910,848 |
| 1999 | 301,957 | 522,201 | 1,024,398 | 4,969,283 | 212,991 | 26,356 | 32,375 | 231,692 | 7,321,253 |
| 2000 | 1,125,730 | 483,963 | 2,672,996 | 4,888,910 | 201,306 | 13,457 | 62,390 | 242,912 | 9,691,664 |
| 2001 | 1,132,214 | 304,127 | 1,278,699 | 7,674,759 | 355,009 | 10,750 | 7,844 | 320,487 | 11,083,889 |
| 2002 | 268,423 | 250,899 | 1,162,278 | 7,075,130 | 242,184 | 29,343 | 10,622 | 117,880 | 9,156,759 |
| 2003 | 682,698 | 262,114 | 2,069,176 | 5,674,111 | 317,606 | 59,399 | 71,881 | 79,396 | 9,216,381 |
| 2004 | 861,987 | 307,898 | 1,078,951 | 5,792,487 | 300,440 | 69,510 | 15,597 | 275,858 | 8,702,728 |
| 2005 | 1,183,631 | 755,232 | 987,748 | 7,240,971 | 163,751 | 34,922 | 14,995 | 145,376 | 10,526,626 |
| 2006 | 638,138 | 729,730 | 864,415 | 6,460,336 | 218,775 | 16,240 | 9,210 | 188,671 | 9,125,515 |
| 2007 | 441,806 | 320,458 | 806,024 | 6,111,612 | 129,675 | 11,109 | 12,756 | 207,030 | 8,040,470 |
| 2008 | 526,458 | 317,997 | 462,531 | 3,612,065 | 133,416 | 16,212 | 12,948 | 320,430 | 5,402,057 |
| 2009 | 127,115 | 239,126 | 1,512,280 | 3,708,788 | 132,895 | 71,517 | 36,771 | 271,949 | 6,100,441 |
| 2010 | 36,087 | 40,166 | 977,562 | 3,185,485 | 233,607 | 12,566 | 10,067 | 109,513 | 4,605,053 |
| 2011 | 21,460 | 52,889 | 443,520 | 1,837,183 | 100,692 | 240,665 | 21,548 | 264,884 | 2,982,841 |
| 2012 | 85,093 | 61,535 | 397,873 | 1,905,100 | 105,541 | 12,291 | 13,503 | 371,635 | 2,952,571 |

Table 4. Recreational harvest (numbers) of Atlantic croaker by state, 1981-2012

(Source: personal communication with NMFS Fisheries Statistics Division, Silver Spring, MD.)

| Year | NJ | DE | MD | VA | NC | SC | GA | FL | Total |
|------|-----------|---------|-----------|-----------|-----------|---------|---------|-----------|------------|
| 1981 | 1,054 | 3,003 | 0 | 964,013 | 1,043,240 | 165,742 | 35,591 | 598,896 | 2,811,539 |
| 1982 | | | 10,452 | 273,039 | 596,493 | 193,554 | 169,749 | 1,682,619 | 2,925,906 |
| 1983 | | | 108,355 | 2,154,133 | 1,620,909 | 60,811 | 75,173 | 1,148,227 | 5,167,608 |
| 1984 | | | 211,035 | 2,047,720 | 2,147,871 | 588,114 | 202,364 | 2,781,742 | 7,978,846 |
| 1985 | | | 21,276 | 2,284,334 | 723,933 | 260,265 | 144,341 | 1,306,955 | 4,741,104 |
| 1986 | | 4,694 | 123,578 | 6,384,966 | 356,742 | 599,442 | 69,887 | 5,118,552 | 12,657,861 |
| 1987 | 0 | 0 | 208,488 | 3,234,224 | 904,030 | 166,978 | 44,783 | 2,580,727 | 7,139,230 |
| 1988 | | 1,186 | 1,005,452 | 4,048,690 | 2,256,128 | 144,057 | 64,093 | 685,778 | 8,205,384 |
| 1989 | | 478 | 22,871 | 2,203,504 | 2,131,763 | 217,023 | 72,598 | 359,417 | 5,007,654 |
| 1990 | | 281 | 100,673 | 2,374,679 | 1,063,452 | 346,631 | 585,380 | 304,064 | 4,775,160 |
| 1991 | 16,235 | 37,500 | 288,471 | 4,298,542 | 434,067 | 100,816 | 184,435 | 1,030,115 | 6,390,181 |
| 1992 | 0 | 9,854 | 117,427 | 4,524,040 | 723,823 | 74,051 | 440,185 | 754,595 | 6,643,975 |
| 1993 | 2,552 | 19,352 | 805,560 | 4,990,098 | 755,998 | 32,700 | 89,734 | 304,067 | 7,000,061 |
| 1994 | 1,567 | 5,718 | 1,633,581 | 6,494,691 | 1,179,735 | 188,520 | 102,974 | 599,032 | 10,205,818 |
| 1995 | 15,184 | 136,865 | 827,183 | 5,029,708 | 850,606 | 75,422 | 100,826 | 438,076 | 7,473,870 |
| 1996 | 35,037 | 235,389 | 775,115 | 4,997,021 | 662,240 | 37,464 | 61,957 | 116,575 | 6,920,798 |
| 1997 | 342,089 | 385,586 | 1,053,232 | 8,066,926 | 661,116 | 118,428 | 64,050 | 235,430 | 10,926,857 |
| 1998 | 143,404 | 391,231 | 1,126,058 | 6,730,181 | 387,427 | 170,528 | 64,953 | 234,360 | 9,248,142 |
| 1999 | 357,261 | 662,724 | 1,209,572 | 5,881,671 | 442,185 | 54,761 | 104,438 | 403,982 | 9,116,594 |
| 2000 | 1,023,442 | 517,886 | 2,674,880 | 5,486,159 | 391,056 | 32,332 | 128,922 | 455,870 | 10,710,547 |
| 2001 | 1,177,813 | 312,005 | 1,319,928 | 9,335,313 | 635,552 | 19,802 | 21,503 | 426,264 | 13,248,180 |
| 2002 | 253,472 | 261,634 | 1,223,385 | 9,129,060 | 408,944 | 66,409 | 36,497 | 177,751 | 11,557,152 |
| 2003 | 692,391 | 341,174 | 1,619,766 | 6,695,192 | 490,399 | 198,339 | 248,853 | 165,459 | 10,451,573 |
| 2004 | 855,927 | 389,218 | 896,855 | 8,259,608 | 511,418 | 171,544 | 38,599 | 415,570 | 11,538,739 |
| 2005 | 1,227,349 | 825,267 | 784,246 | 7,657,147 | 326,777 | 143,387 | 39,561 | 302,784 | 11,306,518 |
| 2006 | 511,220 | 763,216 | 754,969 | 7,221,148 | 556,024 | 58,500 | 34,081 | 172,586 | 10,071,744 |
| 2007 | 406,238 | 359,064 | 872,838 | 6,944,886 | 461,162 | 38,147 | 45,068 | 310,130 | 9,437,533 |
| 2008 | 600,975 | 368,911 | 619,942 | 8,388,497 | 317,940 | 65,853 | 38,246 | 449,054 | 10,849,418 |
| 2009 | 193,464 | 451,849 | 1,335,439 | 5,327,388 | 368,990 | 238,900 | 82,269 | 438,209 | 8,436,508 |
| 2010 | 63,027 | 75,404 | 1,136,589 | 4,743,697 | 478,156 | 46,464 | 35,635 | 132,664 | 6,711,636 |
| 2011 | 40,855 | 92,289 | 554,206 | 3,305,707 | 246,676 | 349,464 | 44,044 | 476,292 | 5,109,533 |
| 2012 | 237,994 | 84,403 | 701,482 | 3,445,232 | 288,812 | 27,541 | 38,402 | 589,643 | 5,413,509 |

Table 5. Recreational releases (number) of Atlantic croaker by state, 1981-2012

(Source: personal communication with NMFS Fisheries Statistics Division, Silver Spring, MD.)

| Year | NJ | DE | MD | VA | NC | SC | GA | FL | Total |
|------|-----------|-----------|-----------|-----------|-----------|---------|---------|-----------|------------|
| 1981 | | | 16,233 | 324,238 | 704,259 | 128,192 | 13,481 | 85,740 | 1,272,143 |
| 1982 | | | | 77,756 | 641,327 | 107,340 | 111,630 | 188,277 | 1,126,330 |
| 1983 | | | 1,507,184 | 1,410,151 | 424,562 | 119,036 | 70,499 | 379,021 | 3,910,453 |
| 1984 | | | 70,192 | 673,080 | 1,701,418 | 746,905 | 37,573 | 236,432 | 3,465,600 |
| 1985 | | | 13,132 | 1,616,052 | 1,596,901 | 238,678 | 66,649 | 1,146,582 | 4,677,994 |
| 1986 | | 1,757 | 43,399 | 2,578,268 | 137,841 | 84,335 | 40,623 | 318,511 | 3,204,734 |
| 1987 | 1,374 | 861 | 32,074 | 2,056,580 | 560,853 | 108,366 | 76,908 | 1,770,697 | 4,607,713 |
| 1988 | | 582 | 273,231 | 832,284 | 984,219 | 112,271 | 20,021 | 200,630 | 2,423,238 |
| 1989 | | 1,307 | 41,822 | 1,342,169 | 891,926 | 58,642 | 17,632 | 72,822 | 2,426,320 |
| 1990 | | 1,268 | 88,688 | 3,922,564 | 1,351,152 | 111,085 | 317,497 | 168,144 | 5,960,398 |
| 1991 | 91,633 | 75,319 | 3,352,190 | 7,418,045 | 669,385 | 25,168 | 140,402 | 647,824 | 12,419,966 |
| 1992 | 4,103 | 43,583 | 856,292 | 4,167,137 | 954,494 | 26,729 | 178,267 | 251,343 | 6,481,948 |
| 1993 | 5,799 | 13,194 | 2,504,362 | 5,795,479 | 1,499,217 | 16,949 | 83,203 | 138,875 | 10,057,078 |
| 1994 | 17,253 | 14,069 | 1,628,824 | 7,676,780 | 3,110,528 | 141,513 | 99,026 | 331,736 | 13,019,729 |
| 1995 | 31,019 | 41,574 | 496,046 | 5,494,289 | 1,172,716 | 108,345 | 89,609 | 141,732 | 7,575,330 |
| 1996 | 17,585 | 76,851 | 403,776 | 5,151,206 | 1,218,799 | 64,494 | 60,282 | 126,300 | 7,119,293 |
| 1997 | 111,468 | 384,233 | 1,497,670 | 7,275,160 | 1,443,568 | 138,107 | 25,630 | 116,276 | 10,992,112 |
| 1998 | 221,324 | 839,932 | 3,021,780 | 4,990,541 | 1,060,928 | 266,068 | 159,928 | 152,744 | 10,713,245 |
| 1999 | 860,325 | 1,017,499 | 2,483,800 | 5,668,925 | 1,368,478 | 116,826 | 57,567 | 967,894 | 12,541,314 |
| 2000 | 688,746 | 694,813 | 4,967,856 | 7,811,048 | 1,569,385 | 96,402 | 169,903 | 428,131 | 16,426,284 |
| 2001 | 853,621 | 285,123 | 1,585,806 | 7,086,706 | 1,256,807 | 115,284 | 192,362 | 282,461 | 11,658,170 |
| 2002 | 369,003 | 361,355 | 2,523,276 | 7,107,656 | 925,806 | 92,498 | 194,474 | 217,054 | 11,791,122 |
| 2003 | 833,508 | 654,697 | 1,393,224 | 6,543,524 | 1,552,315 | 440,446 | 965,496 | 192,356 | 12,575,566 |
| 2004 | 1,237,164 | 599,207 | 854,132 | 6,276,767 | 1,656,049 | 320,788 | 154,259 | 253,951 | 11,352,317 |
| 2005 | 1,692,401 | 674,684 | 1,136,876 | 8,738,109 | 1,401,413 | 321,861 | 280,889 | 293,692 | 14,539,925 |
| 2006 | 503,490 | 937,193 | 1,783,557 | 4,193,675 | 2,578,819 | 595,075 | 283,851 | 187,562 | 11,063,222 |
| 2007 | 590,078 | 672,771 | 1,258,131 | 8,504,212 | 1,608,120 | 224,454 | 228,564 | 321,559 | 13,407,889 |
| 2008 | 2,373,945 | 601,994 | 2,127,219 | 7,806,627 | 1,419,019 | 205,373 | 293,926 | 596,450 | 15,424,553 |
| 2009 | 108,370 | 537,587 | 1,137,578 | 7,621,484 | 1,912,670 | 514,839 | 434,608 | 406,822 | 12,673,958 |
| 2010 | 167,191 | 228,936 | 1,011,236 | 4,824,151 | 1,598,139 | 187,138 | 263,987 | 188,637 | 8,469,415 |
| 2011 | 62,391 | 88,524 | 365,716 | 4,872,928 | 1,798,230 | 240,605 | 262,493 | 452,669 | 8,143,556 |
| 2012 | 1,134,778 | 444,935 | 1,578,524 | 5,091,063 | 1,255,215 | 216,420 | 167,488 | 641,569 | 10,529,992 |