


2013 REVIEW OF THE  
ATLANTIC STATES MARINE FISHERIES COMMISSION  
FISHERY MANAGEMENT PLAN FOR

**SPOTTED SEATROUT**  
*(Cynoscion nebulosus)*

2012 FISHING YEAR



**The Spotted Seatrout Plan Review Team**

Kirby Rootes-Murdy, Atlantic States Marine Fisheries Commission, Chair  
Chip Collier, North Carolina Division of Marine Fisheries  
Chris Kalinowsky, Georgia Coastal Resources Division  
Mike Murphy, Florida Fish & Wildlife Conservation Commission  
Dr. Steve Arnott, South Carolina Department of Natural Resources

September 2013

# 2013 Spotted Seatrout FMP Review

## Table of Contents

I.	Status of the Fishery Management Plan.....	3
II.	Status of the Stock .....	4
III.	Status of the Fishery.....	5
IV.	Status of Assessment Advice .....	6
V.	Status of Research and Monitoring.....	6
VI.	Status of Management Measures and Issues.....	8
VII.	Implementation of FMP Compliance Requirements for 2008.....	8
VIII.	Recommendations of FMP Review Team .....	9
IX.	References.....	9
X.	Figures.....	11
XI.	Tables .....	11

## 2013 Spotted Seatrout FMP Review

### I. Status of the Fishery Management Plan

<u>Date of FMP Approval:</u>	Original FMP – October 1984
<u>Amendments:</u>	Amendment 1 – November 1991 Omnibus Amendment to Spanish Mackerel, Spot, and Spotted Seatrout (Amendment 2)- August 2011
<u>Management Area:</u>	The Atlantic coast distribution of the resource from Maryland through the east coast of Florida
<u>Active Boards/Committees:</u>	South Atlantic State/Federal Fisheries Management Board; Spotted Seatrout Plan Review Team, Omnibus Amendment Plan Development Team

The Atlantic States Marine Fisheries Commission (ASMFC) adopted the Fishery Management Plan (FMP) for spotted seatrout in 1984. The states of Florida through Maryland have a declared interest in the Commission's FMP for spotted seatrout. The ISFMP Policy Board approved Amendment 1 to this FMP in November 1991. In August of 2011, the South Atlantic State/Federal Management Board approved the Omnibus Amendment to Spanish Mackerel, Spot, and Spotted Seatrout FMPs. The Omnibus Amendment (Amendment 2) brought the Spotted Seatrout FMP under the authority of the Atlantic Coastal Fisheries Cooperative Management Act (1993) and the ASMFC Interstate Fishery Management Plan Charter (1995).

The goal of the management plan is "to perpetuate the spotted seatrout resource in fishable abundance throughout its range and generate the greatest possible economic and social benefits from its harvest and utilization over time." Plan objectives include: 1) attain over time optimum yield; 2) maintain a spawning potential ratio of at least 20% to minimize the possibility of recruitment failure; 3) promote conservation of the stocks in order to reduce the inter-annual variation in availability and increase yield per recruit; 4) promote the collection of economic, social, and biological data required to effectively monitor and assess management efforts relative to the overall goal; 5) promote research that improves understanding of the biology and fisheries of spotted seatrout; 6) promote harmonious use of the resource among various components of the fishery through coordination of management efforts among the various political entities having jurisdiction over the spotted seatrout resource; and 7) promote determination and adoption of standards of environmental quality and provide habitat protection necessary for the maximum natural protection of spotted seatrout. Amendment 2 to the Spotted Seatrout FMP added the following objectives in support of the compliance under the Act: 1) Manage the spotted seatrout fishery restricting catch to mature individuals; 2) manage the spotted seatrout stock to maintain sufficiently-high spawning stock biomass; 3) develop research priorities that will further refine the spotted seatrout management program to maximize the biological, social, and economic benefits derived from the spotted seatrout population.

Recommended management measures include a minimum size limit of 12 inches total length (TL) with comparable mesh size regulations in directed fisheries, and data collection for stock assessment and monitoring the status of the fisheries. All states with a declared interest in spotted seatrout have implemented at least the recommended minimum size limit. In addition, each state

has either initiated spotted seatrout data collection programs or modified other programs to collect improved catch and effort data. Table 1 provides the states' recreational and commercial regulations for spotted seatrout in 2012.

## II. Status of the Stock

A coastwide stock assessment of spotted seatrout has not been conducted given the largely non-migratory nature of the species and the lack of data on migration where it does occur. Instead, state-specific age-structured analyses of local stocks have been performed by several states. These stock assessments provide estimates of spawning potential ratio (SPR), which is a measure of the effect of fishing pressure on the relative abundance of the mature female segment of the population. The FMP recommends a goal of 20% SPR; North Carolina, South Carolina, and Georgia have adopted this goal, and Florida has established a 35% SPR goal.

Florida's stock assessments are for separate northern and southern populations. Average transitional SPR estimates for Florida's spotted seatrout during 2007-2009 were 67% in the northeast region of the state's Atlantic coast and 45% in the southeast region (Murphy et al. 2011). This assessment provided the basis for some relaxation in the management of spotted seatrout in Florida (Table 1).

The South Carolina Department of Natural Resources packaged three state-specific assessments into a report in 2001; however, these assessments were not peer reviewed. This initial assessment of South Carolina spotted seatrout covered 1986-1992 and indicated that female SPR was just above the 20% goal in the terminal year (Zhao and Wenner 2001). This assessment led to an increase in the minimum size limit and decrease in the creel limit for spotted seatrout in South Carolina. A more recent assessment of the population of South Carolina spotted seatrout was conducted for the period 1981-2004, but not peer reviewed (de Silva, Draft 2005). Two modeling approaches were used, and both models indicated that the current spawning stock biomass is below what would be required to maintain 20% SPR.

Assessments in North Carolina and Georgia spotted seatrout covered 1981-1997 and 1986-1995, respectively, and both indicated that female SPR was below the 20% goal in the terminal year (Zhao and Burns 2001, Zhao *et al.* 2001). A more recent assessment of spotted seatrout in Georgia has been performed; however, it remains unpublished. This 2002 Georgia assessment is unpublished because the results were highly questionable due to data deficiencies and changing methodologies.

North Carolina recently completed a peer reviewed stock assessment of spotted seatrout covering 1991-2008, which included all spotted seatrout caught in North Carolina and Virginia (Jensen 2009). The assessment indicated that SPR has been below 20% SPR in recent years. Jensen (2009) recommended the implementation of management measures to account for recent increases of recreational fishing and discard mortality and maintain a sufficiently large spotted seatrout population to act as a buffer against the effects of future cold stun events. Based on the assessment, North Carolina developed a draft state FMP for spotted seatrout, with the final version approved in April 2012.

### **III. Status of the Fishery**

Both commercial and recreational fishermen regularly catch spotted seatrout from Maryland through the east coast of Florida (except in South Carolina where spotted seatrout has been declared a gamefish and can only be taken by recreational means). Landings from states north of Maryland are minimal and/or inconsistent from year to year. All catch estimates in this section include those in the management area only (MD-FL). Total recreational landings have surpassed total commercial landings every year since recreational landings have been recorded in 1981 (Figure 1). In 2009, recreational landings were more than five times the commercial landings. A coastwide (VA, NC, SC) winter mortality event in 2000/2001 likely contributed to the sudden decline in commercial and recreational landings in 2001 and 2002. Both fisheries' landings have increased since then.

#### *Commercial Fishery*

The National Marine Fisheries Service (NMFS) compiles commercial spotted seatrout landings. The data are cooperatively collected by the NMFS and state fishery agencies from state mandated trip-tickets, landing weigh-out reports from seafood dealers, federal logbooks, shipboard and portside interviews, and biological sampling of catches. See Table 2.

Atlantic coast commercial landings of spotted seatrout (1960-2012) have ranged from 165,000 pounds to 1.38 million pounds (Figure 1). Commercial landings historically came primarily from Florida and North Carolina, with Virginia, South Carolina, and Georgia accounting for a small portion of the total. From 1960 to 1976, annual commercial landings of spotted seatrout averaged 1.07 million pounds, but have declined since then due to increased regulation and possible declines in abundance. Significant changes to regulations include the 1987 designation of spotted seatrout as a gamefish in South Carolina, and the 1995 prohibition on the use of entangling nets in Florida's coastal waters. From 2002 to 2011, commercial landings have averaged approximately 292,022 pounds. North of Florida, variability in annual harvest is typical and seems to parallel the climatic conditions of the preceding winter and spring. In 2012 the commercial landings are estimated to be 408,520 pounds, representing a 161% increase from the previous year's harvest and a 39.9% increase from the previous ten-year average. North Carolina accounted for approximately 65% of the total coastwide catch, with Virginia and Florida responsible for approximately 19% and 15% of the 2012 commercial landings, respectively.

#### *Recreational Fishery*

Recreational catch statistics are collected by the NMFS recreational fisheries survey. Effort data are collected through telephone interviews. Catch data are collected through access-point angler intercept surveys. Catch per trip estimates are produced for each type of fish encountered, either observed or reported, and these estimates are combined with the effort estimates by sampling stratum to produce the catch and harvest estimates. See Tables 3, 4, and 5.

Over the last 28 years, the recreational catch of spotted seatrout (kept and released) has shown a strong upward trend, increasing from 1.1 million fish in 1981 to a peak of 8.8 million fish in 2012 (Figure 2). The recreational harvest of spotted seatrout, however, has remained relatively stable around the time series average of 1.3 million fish. The recreational harvest increased from approximately 952,458 fish in 2010 to 1.8 million in 2012. Due in part to recreational size and creel limits and closed seasons, as well as the encouragement of catch and release practices, the

percentage of caught fish being released has increased to 75-87 percent of the catch since 2000. In 2012, the release percentage (79.4%) was similar to the previous 10-year average (76.3%). In 2012, Georgia anglers took the largest proportion of harvested fish with 29%, followed closely by North Carolina anglers at 27%. Recreational catches are generally made with rod and reel, but some are taken by recreational nets and by gigging, where these methods are permitted. Most recreational fishing is conducted from private boats and the majority of the catch is taken from nearshore waters.

#### **IV. Status of Assessment Advice**

A coastwide stock assessment of spotted seatrout has not been conducted and the Plan Review Team (PRT) does not recommend that one be completed due to the life history of the fish and the available data. Several states have performed age-structured analyses on local stocks of spotted seatrout. Recent Florida and North Carolina stock assessments for spotted seatrout provide divergent trends on the status of the species. The 2005 stock assessment in South Carolina indicated an increasing population trend but a status level that is still below target spawning stock biomass levels (de Silva 2005). The PRT supports the continuation of state-specific assessments, yet recognizes the difficulty most states face to attain sufficient data of a quality that can be used in the assessment process and personnel who can perform the necessary modeling exercises.

The lack of biological and fisheries data for stock assessment and effective management of the resource was recognized in the 1984 FMP and continues to be a hindrance. Some states are increasing their collection of biological and fisheries data, which should provide insight on stock status over time.

#### **V. Status of Research and Monitoring**

In addition to the commercial and recreational fishery-dependent data collected and/or compiled through the National Marine Fisheries Service, Fisheries Statistics Division, some states have implemented fishery-independent or additional fishery-dependent monitoring programs.

The Florida Fish and Wildlife Conservation Commission (FWC) implemented a juvenile finfish monitoring program in the northern Indian River Lagoon in the spring of 1990 and in the estuarine reaches of the St. Johns, St. Marys, and Nassau Rivers in northeast Florida in the spring of 2001 (FWC-FWRI 2013). Florida also initiated a stratified random sampling program in 1997 on the Atlantic coast that utilizes a 183-m haul seine to catch exploitable-sized fishes. This has been conducted in the northern Indian River and southern Indian River since initiation and in northeast Florida since 2001. Trends in the YOY abundance have been relatively stable with periods of strong recruitment evident. Recent strong recruitment appears to have occurred in Northeast Florida (2011 and 2012) but is not as evident in the central and southern areas of Florida's Atlantic coast. For 2012 sampling program, 191 lengths were measured and 131 otoliths were collected from adult sized spotted seatrout.

Florida's fishery-dependent sampling includes commercial trip-ticket information and biostatistical sampling of the commercial and recreational catch. A voluntary angler logbook program was implemented in 2002 to collect information on the lengths of spotted seatrout

## 2013 Spotted Seatrout FMP Review

released alive by anglers. Recently (2011) this program changed to 'postcard' program enlisting anglers encountered at sites visited during the MRIP angler intercept survey.

Georgia collects fishery-dependent data through a Marine Sportfish Carcass Recovery Program. Data collected through this survey are used to examine trends in the size and age composition of the recreationally harvested population, valuable information for future stock assessments. For 2012, a total of 4,428 fish carcasses were donated through the program. Approximately 77% (3,431) of the carcasses were seatrout, with an average centerline (CL) length of 362.2 mm CL (minimum: 290 mm CL; maximum: 651 mm CL), were reported from 16 recovery locations.

Georgia also collects fishery-independent data through the Marine Sportfish Population Health Study, was implemented in 2003 to provide age and sex specific estimates of relative abundance in two Georgia estuaries, Wassaw Sound and the Altamaha Sound region. This trammel net survey is conducted monthly, September through November, and utilizes a hybrid random-stratified and fixed station design in which each station is sampled once in a given month. For 2012, the average centerline length in Wassaw was 362.4 mm CL and 351.0 mm CL in Altamaha.

South Carolina has an extensive directed research program on this species. Current project objectives include determining the size and age composition of the recreational catch by sampling independent angler and fishing tournament catches as well as a carcass program, and producing fisheries independent relative abundance estimates from trammel net surveys along the South Carolina coast. The latter is a stratified random sampling design and has been conducted monthly since November 1990. South Carolina also has an electrofishing survey of upper estuarine waters. It uses a stratified random design and has been operating monthly since 2001. In 2012, a total of 50 Spotted Seatrout were captured by random electrofishing sets, with a mean overall CPUE of 0.24 Spotted Seatrout per set. CPUE has generally declined in the electrofishing survey since 2009. In contrast to electrofishing, the trammel net survey, catches some YOY as well as older seatrout (S. Arnott, Personal Communication, 2011). During 2012, a total of 3,258 Spotted Seatrout were captured in random trammel net sets, with an overall mean CPUE of 3.37 Spotted Seatrout per trammel set. Additionally, South Carolina also has ongoing seatrout parasite studies (Moravec et al. 2006). Catch rates, size composition, and sub-samples of the catch on a bi-monthly basis are used for generating age-length keys for cohort specific indices of abundance. Roumillat and Brouwer (2004) have described the reproductive dynamics of female spotted seatrout in South Carolina.

North Carolina has collected age, growth, and maturity data for spotted seatrout caught in fishery-dependent and fishery-independent sampling programs since 1991. A fishery-independent monitoring program was initiated in May 2001, supported by USFWS Sports Fish Restoration funds. The program utilizes a stratified random, multi-mesh size gill net survey along North Carolina's Outer Banks, the bays of western Pamlico Sound, the Neuse, the Pamlico, Pungo, New and Cape Fear Rivers, and the Atlantic Ocean. Project objectives include calculating annual indices of abundance for important recreational fish (spotted seatrout included); supplementing samples for age, growth, and reproductive studies; evaluating catch rates and species distribution for identifying and resolving bycatch problems; and characterizing habitat utilization. Additional areas of the Neuse and Pamlico-Pungo Rivers contribute to the

## 2013 Spotted Seatrout FMP Review

Pamlico Sound Area Independent Gill Net Survey, with common objectives and sampling design. Hydrophone work was conducted in North Carolina to characterize critical spawning habitats for spotted seatrout in Pamlico Sound. For the 2012 surveying program, the overall spotted seatrout CPUE was 0.68 (n=193) for Pamlico Sound (second highest in the time series); 0.64 (n=204) for surveys in the Pamlico-Pungo, and Neuse rivers (also second highest in the time series); and 0.37 (n=45) for surveys in the Cape Fear and New Rivers. Hook-and-line and estuarine gill net discard mortality studies were conducted in North Carolina in 1998-2001, supported by Atlantic Coastal Fisheries Cooperative Management Act funds.

A spotted seatrout tagging study was initiated in September 2008 and is scheduled to conclude in August 2012. Funding for one year was to collect preliminary data necessary to design and conduct an effective long-term tagging study on spotted seatrout in North Carolina, 2008-2009 (funded by NC Sea Grant Fishery Resource Grant). This was followed by an advanced tagging study by NC State University researchers who are using a combined conventional tag and telemetry approach to study the movement and mortality of spotted seatrout in North Carolina, 2009-2012 (funded by NC Marine Resources Fund, which consists of proceeds from the sale of the Coastal Recreational Fishing License).

### **VI. Status of Management Measures and Issues**

#### *Changes to State Regulations*

##### North Carolina:

Reduction in recreational bag limit from six fish to four fish and removed restriction limiting two fish to greater than 24 inches total length.

##### Florida

Effective September 1, 2013, the recreational seasons were dropped, the commercial season was lengthened, and the commercial possession limit was modified to accommodate twice the possession limit on a vessel occupied by two or more license fishers.

#### *Omnibus Amendment (Interstate)*

In August 2011, the Management Board approved an amendment to the Spotted Seatrout FMP to address three issues: compliance measures, consistency with federal management in the exclusive economic zone, and alignment with Commission standards. The updated FMP's objectives are to: (1.) Manage the spotted seatrout fishery restricting catch to mature individuals. (2.) Manage the spotted seatrout stock to maintain sufficiently-high spawning stock biomass. (3.) Develop research priorities that will further refine the spotted seatrout management program to maximize the biological, social, and economic benefits derived from the spot population. Through the Omnibus Amendment requires the following fishery management measures in either the recreational or commercial fisheries for states within the management unit range:

12"TL minimum size with comparable mesh size requirements

#### De minimis Guidelines



A state qualifies for *de minimis* status if its past 3-years' average of the combined commercial and recreational catch is less than 1% of the past 3-years' average of the coastwide combined commercial and recreational catch. Those states that qualify for *de minimis* are not required to implement any monitoring requirements, none of which are included in the plan.

### **De Minimis Requests**

The state of New Jersey requests *de minimis* status. The PRT notes they meet the requirements of *de minimis*.

## **VII. Implementation of FMP Compliance Requirements for 2012**

12'' TL minimum size with comparable mesh size requirements (both commercial and recreational)

## **VIII. Recommendations of Plan Review Team**

### Management and Regulatory Recommendations

- Increase observer coverage in states that have a commercial fishery for spotted seatrout.

### Prioritized Research Recommendations

#### *High Priority*

- Conduct state-specific stock assessments to determine the status of stocks relative to the plan objective of maintaining a spawning potential of at least 20%.
- Collect data on the size or age of spotted seatrout released alive by anglers and the size and age of commercial discards.
- Continue work to examine the stock structure of spotted seatrout on a regional basis, with particular emphasis on advanced tagging techniques.
- Expand the NMFS recreational fishery survey to assure adequate data collection for catch and effort data, increased intercepts, and state add-ons of social and economic data needs.
- Conduct telemetry tagging surveys to provide precise estimates of mortality attributed to winter kills.
- Provide state-specific batch fecundity estimates for use in stock assessments.
- Develop state-specific juvenile abundance indices.
- Increase observer coverage in states that have a commercial fishery for spotted seatrout.

#### *Medium Priority*

- Identify essential habitat requirements.
- Evaluate effects of environmental factors on spawning frequency and stock density.
- Initiate collection of social and economic aspects of the spotted seatrout fishery.

## **IX. References**

de Silva, JA. 2005. Draft. Stock assessment of spotted seatrout, *Cynoscion nebulosus*, in South Carolina with recommendations on the management of the recreational fishery. South Carolina Department of Natural Resources, Marine Research Institute, Charleston, South Carolina.


Florida Fish and Wildlife Conservation Commission's Fish and Wildlife Research Institute

## 2013 Spotted Seatrout FMP Review


- (FWC-FWRI). 2013. Species Profile: Spotted Seatrout. *In*: R.H. McMichael, editor) Fisheries-independent monitoring program, 2012 annual data summary report, St. Petersburg.
- Jensen CC. 2009. Stock Status of Spotted Seatrout, *Cynoscion nebulosus*, in North Carolina, 1991-2008. Morehead City (NC): North Carolina Division of Marine Fisheries. 89 p.
- Moravec F, de Buron I, Roumillat WA. 2006. Two new species of Philometra (Nematoda : Philometridae) parasitic in the perciform fish *Cynoscion nebulosus* (Sciaenidae) in the estuaries of South Carolina, USA. *Folia Parasitologica*, 53: 63-70
- Murphy MD, D.Chagaris, and D. Addis. 2011. An assessment of the status of spotted seatrout in Florida waters through 2009. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute. In-House Report 2011-002, St. Petersburg.
- Roumillat WA, Brouwer MC. 2004. Reproductive dynamics of female spotted seatrout (*Cynoscion nebulosus*) in South Carolina. *Fisheries Bulletin*, 102: 473-487
- Zhao B, Burns B. 2001. Stock assessment of the spotted seatrout, *Cynoscion nebulosus*, on the North Carolina Coast, 1981-1997. *In*: South Carolina Department of Natural Resources. Cooperative Research on the Biology and Assessment of Nearshore and Estuarine Fishes along the Southeast Coast of the U.S: Part III. Spotted Seatrout, *Cynoscion nebulosus*. Charleston (SC): SC DNR. Final Report, Grant NA77FF0550.
- Zhao B, Wenner C. 2001. Stock assessment of the spotted seatrout, *Cynoscion nebulosus*, on the South Carolina Coast, 1986-1992. *In*: South Carolina Department of Natural Resources. Cooperative Research on the Biology and Assessment of Nearshore and Estuarine Fishes along the Southeast Coast of the U.S: Part III. Spotted Seatrout, *Cynoscion nebulosus*. Charleston (SC): SC DNR. Final Report, Grant NA77FF0550.
- Zhao B, Wenner C, Nicholson N. 2001. Stock assessment of the spotted seatrout, *Cynoscion nebulosus*, on the Georgia Coast, 1986-1995. *In*: South Carolina Department of Natural Resources. Cooperative Research on the Biology and Assessment of Nearshore and Estuarine Fishes along the Southeast Coast of the U.S: Part III. Spotted Seatrout, *Cynoscion nebulosus*. Charleston (SC): SC DNR. Final Report, Grant NA77FF0550.

**X. Figures**

**Figure 1. Commercial landings (1960-2012) and recreational landings (1981-2012), in pounds, from Maryland to Florida (See Tables 2 and 4 for values and sources)**



**Figure 2. Recreational catch (numbers), 1981-2012, from Maryland to Florida (See Tables 3 and 5 for values and sources)**



2013 Spotted Seatrout FMP Review

**XI. Tables**

**Table 1. Summary of state regulations for spotted seatrout in 2012**

State	Recreational	Commercial
New Jersey	13" TL; 1 fish	Gill net: 13"; open 1/1-5/20 & 9/3-10/19 & 10/27-12/31; 100 lb possession limit; 100 lb bycatch limit; mesh $\geq 3.25$ " stretched except 2.75 - 3.25" stretched allowed within 2nm for permitted fishermen doing monthly reporting. Trawl: 13"; open 1/1-7/31 & 10/13-12/31; mesh $\geq 3.75$ " diamond or 3.375 square; 100 lb possession limit' 100 lb bycatch limit. Pound net: 13"; open 1/1/-6/6 & 7/1-12/31; 150 lb bycatch limit. Hook & line: open 1/1-12/31, 13", 1 fish.
Delaware	12" TL	12" TL. Gill net restrictions.
Maryland	14" TL; 10 fish	12" TL. Minimum mesh size restrictions for trawl (3-3/8" sq. or 3-3/4" diag.) and gill nets (3").
PRFC	14" TL; 10 fish	14" TL
Virginia	14" TL; 10 fish	14" TL except pound nets and haul seines allowed 5% by weight less than 14". Hook & line - 10 fish limit. Quota: 51,104 lbs (Sept. 1-Aug. 31).
North Carolina	14" TL; 4 fish	14" TL; hook & line - 75 fish limit.
South Carolina	14" TL; 10 fish. May be taken by rod & reel year-round or gigging March-November.	Gamefish status: native caught fish may not be sold.
Georgia	13" TL; 15 fish	13" TL; 15 fish. BRD requirement for trawl; gear mesh regulations.
Florida	15-20" TL slot with 1 fish >20" allowed; north region: 6 fish limit; south region: 4 fish limit	15-24" TL; June 1-Nov 30 season (north), May 1-Sept 30 season (south); 75 fish per day or vessel (up to 150 per day if two or more licensed commercial fishers aboard); hook & line or cast net only

Note: A commercial fishing license is required to possess spotted seatrout in all states with a fishery.

2013 Spotted Seatrout FMP Review

**Table 2. Commercial landings (pounds) of spotted seatrout by state, 1981-2012**  
(Source: NMFS Fisheries Statistics Division, 09/30/2013)

<b>Year</b>	<b>MD</b>	<b>VA</b>	<b>NC</b>	<b>SC</b>	<b>GA</b>	<b>FL</b>	<b>Total</b>
1981		4,000	113,304		629	736,026	853,959
1982		3,400	83,847	1,944	4,994	732,278	826,463
1983		4,400	165,360	4,479	5,795	481,535	661,569
1984		3,000	152,934	2,374	4,348	367,541	530,197
1985		8,302	109,048	1,770	7,149	369,756	496,025
1986		18,500	191,514	12,214	8,691	307,261	538,180
1987		13,300	315,380	11,941	10,739	317,044	668,404
1988		15,500	296,538	486	9,110	315,947	637,581
1989		18,500	451,909	33	10,565	361,973	842,980
1990		21,435	250,634	1,095	5,942	236,453	515,559
1991	98	21,200	660,662		7,380	225,812	915,152
1992	364	10,395	526,271		11,310	247,189	795,529
1993	24	38,033	449,886		8,550	223,931	720,424
1994	30	44,636	412,458		5,112	247,666	709,902
1995	182	28,722	574,410		8,482	184,269	796,065
1996	14,961	4,476	226,668		7,501	48,254	301,860
1997	15,688	11,711	232,583		7,621	57,316	324,919
1998	19,794	21,774	307,777		2,845	41,556	393,746
1999	36,365	38,513	546,775		3,244	61,802	686,699
2000	20,270	19,918	376,657		1,997	45,392	464,234
2001	24,754	3,773	105,797			30,234	164,558
2002	11,771	9,308	175,643		969	44,640	242,331
2003	902	5,310	181,529			27,075	215,676
2004	342	17,290	131,019		815	29,605	187,700
2005	2,410	21,448	129,645			36,762	210,280
2006	245	28,529	312,714			36,687	398,897
2007	32	41,003	374,817	0	0	46,838	476,804
2008	0	43,666	304,504	0	0	20,887	369,057
2009	243	27,762	320,336	0	0	46,297	394,695
2010		28,346	200,562			39,374	268,282
2011	557	17,107	75,239			63,592	156,495
2012	1890	79,490	265,476	0	0	61,664	408,520

2013 Spotted Seatrout FMP Review

**Table 3. Recreational harvest (numbers of fish) of spotted seatrout by state, 1981-2012**  
(Source: NMFS Fisheries Statistics Division, 09/30/2013)

Year	MD	VA	NC	SC	GA	FL	Total
1981			30,037	20,934	189,080	576,847	816,898
1982			112,023	849,634	226,758	426,378	1,614,793
1983			91,956	121,940	325,655	645,120	1,184,671
1984			90,262	95,281	114,403	700,876	1,000,822
1985			263,878	347,851	251,764	866,162	1,729,655
1986	7,507	82,671	270,867	477,136	401,490	550,591	1,790,262
1987	29,295	17,415	320,977	392,329	439,782	744,330	1,944,128
1988	20,769	288,705	420,115	355,547	389,276	331,709	1,806,121
1989	151,986	66,033	181,149	174,011	448,767	198,617	1,220,563
1990	20,416	67,939	251,088	113,160	368,787	249,824	1,071,214
1991	17,995	69,032	316,895	438,502	1,204,116	385,817	2,432,357
1992	3,235	30,091	333,990	200,030	338,175	363,238	1,268,759
1993	7,038	103,131	206,523	222,144	463,702	274,118	1,276,656
1994	33,511	115,025	457,636	139,551	337,965	255,216	1,338,904
1995	19,198	90,838	325,927	223,751	607,095	381,884	1,648,693
1996	35,765	46,098	151,380	137,530	171,676	148,571	691,020
1997	19,951	92,725	256,719	111,576	167,287	228,096	876,354
1998	13,620	34,623	294,501	125,038	197,293	189,621	854,696
1999	2,112	138,492	410,321	101,260	655,407	241,096	1,548,688
2000	1,634	90,135	250,450	219,740	486,673	288,443	1,337,075
2001		13,447	182,124	63,452	309,487	250,987	819,497
2002		16,303	197,484	84,777	271,357	206,310	776,231
2003	2,091	102,484	106,415	123,027	425,993	169,587	929,597
2004		74,747	316,894	247,156	336,254	199,523	1,174,574
2005	3,828	31,416	512,262	268,467	231,429	337,744	1,385,146
2006	5,136	56,475	577,537	294,096	453,394	299,337	1,685,975
2007		145,736	525,156	122,419	499,709	302,625	1,595,645
2008		79,545	584,024	175,975	623,619	160,455	1,623,618
2009	11,680	40,109	509,416	147,266	478,895	182,752	1,370,118
2010	3,146	17,417	195,065	101,053	384,077	251,455	952,213
2011	3,058	247,736	215,922	66,207	289,950	286,501	1,109,374
2012	6,032	125,627	500,522	234,921	526,604	427,469	1,821,175

2013 Spotted Seatrout FMP Review

**Table 4. Recreational harvest (pounds of fish) of spotted seatrout by state, 1981-2012**  
(Source: NMFS Fisheries Statistics Division, 09/30/2013)

Year	MD	VA	NC	SC	GA	FL	Total
1981			63,036	14,808	138,720	967,921	1,184,485
1982			120,045	588,999	177,847	660,295	1,547,186
1983			96,359	138,442	323,889	784,531	1,343,221
1984			39,861	116,118	141,306	866,077	1,163,362
1985			288,088	509,551	234,704	1,032,344	2,064,687
1986	4,960	64,394	328,439	587,570	440,774	695,168	2,121,305
1987	22,511	38,495	366,442	592,612	491,317	883,707	2,395,084
1988	36,629	460,378	390,836	448,473	536,959	453,063	2,326,338
1989	184,318	112,344	259,726	277,489	608,009	328,338	1,770,224
1990	39,059	121,136	282,872	174,845	423,815	475,045	1,516,772
1991	34,753	121,604	472,397	628,011	1,449,853	534,371	3,240,989
1992	7,802	56,685	508,760	227,210	430,946	543,491	1,774,894
1993	12,800	201,562	307,151	268,055	586,426	392,827	1,768,821
1994	26,764	175,184	679,996	183,343	412,392	357,441	1,835,120
1995	31,464	148,544	478,674	247,987	667,379	642,670	2,216,718
1996		77,269	197,261	171,727	196,487	249,898	892,642
1997	32,963	261,911	311,891	163,771	242,506	380,276	1,393,318
1998	37,189	61,888	444,441	151,718	262,896	329,793	1,287,925
1999		290,694	690,606	146,277	916,860	428,061	2,472,498
2000	2,972	195,544	385,190	267,297	565,903	545,202	1,962,108
2001		26,733	213,438	58,885	369,083	502,254	1,170,393
2002		28,882	274,100	111,954	302,559	353,693	1,071,188
2003	3,494	218,061	145,936	140,276	502,278	316,279	1,326,324
2004		134,602	385,624	229,541	377,370	390,880	1,518,017
2005	10,761	76,325	628,739	326,501	263,209	603,891	1,909,426
2006	9,993	132,629	941,161	369,165	531,441	533,121	2,517,510
2007		305,599	988,527	211,225	531,637	594,506	2,631,494
2008		195,987	922,733	302,019	733,307	298,679	2,452,725
2009	13,261	85,358	833,568	199,554	579,270	322,941	2,033,952
2010	6,724	28,146	407,193	138,514	425,854	411,495	1,418,046
2011	4,664	549,976	403,160	116,979	353,472	464,863	1,893,114
2012	10,257	226,556	817,451	388,105	518,189	819,009	2,779,567

2013 Spotted Seatrout FMP Review

**Table 5. Recreational releases (number of fish) of spotted seatrout by state, 1981-2012**  
(Source: NMFS Fisheries Statistics Division, 09/30/2013)

Year	MD	VA	NC	SC	GA	FL	Total
1981				5,522	36,853	209,059	251,434
1982				8,007	17,645	171,093	196,745
1983			16,579	32,860	12,038	367,881	429,358
1984			30,173	44,436	16,174	76,346	167,129
1985			16,578	6,409	22,917	66,960	112,864
1986	13,639	28,606	19,792	115,315	189,798	35,646	402,796
1987		30,070	136,104	130,253	176,415	41,391	514,233
1988	26,999	148,934	74,818	78,568	182,628	431,665	943,612
1989	52,859	11,977	82,909	54,279	167,025	187,406	556,455
1990	4,874	23,435	84,235	35,223	114,624	203,439	465,830
1991	21,811	40,550	169,921	51,415	369,972	789,779	1,443,448
1992	701	19,855	139,616	97,813	192,261	597,254	1,047,500
1993		65,605	149,744	92,101	146,665	780,573	1,234,688
1994	32,466	243,463	207,262	220,941	125,421	574,629	1,404,182
1995	157,530	327,643	277,896	194,996	327,835	1,074,703	2,360,603
1996	51,594	165,169	153,051	107,691	63,585	1,081,893	1,622,983
1997	4,826	168,964	98,377	89,147	61,148	1,449,278	1,871,740
1998	49,460	74,569	73,024	151,935	100,059	1,005,443	1,454,490
1999	7,082	152,120	253,442	92,792	160,801	1,577,378	2,243,615
2000	4,805	264,550	90,070	368,332	547,765	2,310,491	3,586,013
2001		110,308	194,982	38,709	365,140	1,995,635	2,704,774
2002		136,265	385,162	147,962	357,953	2,326,420	3,353,762
2003		207,270	131,619	314,642	737,730	1,707,957	3,099,218
2004	9,430	295,518	300,025	333,537	608,193	1,969,884	3,516,587
2005	4,612	277,307	817,036	395,483	678,057	3,446,336	5,618,831
2006	9,721	125,135	559,786	666,865	872,395	2,889,495	5,123,397
2007	2,231	414,709	973,516	560,272	957,682	3,623,247	6,531,657
2008		373,146	1,005,298	850,006	719,622	2,140,752	5,088,824
2009	30,381	171,028	1,213,526	398,971	915,301	1,641,702	4,370,909
2010	107,017	550,118	1,684,872	407,228	742,215	2,937,411	6,428,861
2011	7,685	1,214,620	1,916,249	279,969	552,123	2,141,212	6,111,858
2012	55,183	428,540	1,646,512	817,017	1,029,479	3,025,556	7,002,287