

Research Priorities and Recommendations to Support Interjurisdictional Fisheries Management

ATLANTIC CROAKER

Recommendations from the 2017 Atlantic Croaker Benchmark Stock Assessment

Short-term

HIGH PRIORITY

- Increased observer coverage for commercial discards, particularly the shrimp trawl fishery. Develop a standardized, representative sampling protocol for collection of individual lengths and ages of discarded finfish.
- Describe the coast-wide distribution, behavior, and movement of croaker by age, length, and season, with emphasis on collecting larger, older fish.

MEDIUM PRIORITY

- Conduct studies of discard mortality for recreational and commercial fisheries by each gear type in regions where removals are highest.
- In the recreational fishery, develop sampling protocol for collecting lengths of discarded finfish and collect otolith age samples from retained fish.
- Encourage fishery-dependent biological sampling, with proportional landings representative of the distribution of the fisheries. Develop and communicate clear protocols on truly representative sampling.

Long-term

HIGH PRIORITY

- Continue state and multi-state fisheries-independent surveys throughout the species range and subsample for individual lengths and ages. Ensure NEFSC trawl survey continues to take lengths and ages. Examine potential factors affecting catchability in long-term fishery independent surveys.
- Quantify effects of BRDs and TEDs implementation in the shrimp trawl fishery by examining their relative catch reduction rates on Atlantic croaker.
- Continue to develop estimates of length-at-maturity and year-round reproductive dynamics throughout the species range. Assess whether temporal and/or density-dependent shifts in reproductive dynamics have occurred.
- Re-examine historical ichthyoplankton studies for an indication of the magnitude of estuarine and coastal spawning. Pursue specific estuarine data sets from the states (NJ, VA, NC, SC, DE, ME) and coastal data sets (MARMAP, EcoMon).

MEDIUM PRIORITY

- Investigate environmental covariates in stock assessment models, including climate cycles (e.g., Atlantic Multi-decadal Oscillation, AMO, and El Nino Southern Oscillation, El Nino) and

recruitment and/or year class strength, spawning stock biomass, stock distribution, maturity schedules, and habitat degradation.

- Use NMFS Ecosystem Indicators bi-annual reports to consider folding indicators into the assessment; identify mechanisms for how environmental indicators affect the stock
- Encourage efforts to recover historical landings data, determine whether they are available at a finer scale for the earliest years than are currently reported.
- Collect data to develop gear-specific fishing effort estimates and investigate methods to develop historical estimates of effort.
- Develop gear selectivity studies for commercial fisheries with emphasis on age 1+ fish.
- Conduct studies to measure female reproductive output at size and age (fecundity, egg and larval quality) and impact on assessment models and biomass reference points
- Develop and implement sampling programs for state-specific commercial scrap and bait fisheries in order to monitor the relative importance of Atlantic croaker. Incorporate biological data collection into program.
- Investigate the relationship between estuarine nursery areas and their proportional contribution to adult biomass. I.e., are select nursery areas along Atlantic coast ultimately contributing more to SSB than others, reflecting better quality juvenile habitat?