

Research Priorities and Recommendations to Support Interjurisdictional Fisheries Management

ATLANTIC STRIPED BASS

(Full citation: Northeast Fisheries Science Center (NEFSC). 2019. 66th Northeast Regional Stock Assessment Workshop (66th SAW) Assessment Report. US Dept. Commer. Northeast Fish. Sci. Cent. Ref. Doc. 19-08; 1170 p.)

Fishery-Dependent Priorities

High

- Continue collection of paired scale and otolith samples, particularly from larger striped bass, to facilitate development of otolith-based age-length keys and scale-otolith conversion matrices.
- Develop studies to provide information on gear specific (including recreational fishery) discard mortality rates and to determine the magnitude of bycatch mortality.
- Conduct study to directly estimate commercial discards in the Chesapeake Bay.
- Collect sex ratio information on the catch and improve methods for determining population sex ratio for use in estimates of female SSB and biological reference points.

Moderate

- Improve estimates of striped bass harvest removals in coastal areas during wave 1 and in inland waters of all jurisdictions year round.

Fishery-Independent Priorities

High

- Develop and index of relative abundance from the Hudson River Spawning Stock Biomass survey to better characterize the Delaware Bay/Hudson River stock.
- Improve the design of existing spawning stock surveys for Chesapeake Bay and Delaware Bay.

Moderate

- Develop a refined and cost-efficient, fisheries-independent coastal population index for striped bass stocks.
- Collect sex ratio information from fishery-independent sources to better characterize the population sex ratio.

Modeling / Quantitative Priorities

High

- Develop better estimates of tag reporting rates; for example, through a coastwide tagging study.
- Investigate changes in tag quality and potential impacts on reporting rate.

- Explore methods for combining tag results from programs releasing fish from different areas on different dates.
- Develop field or modeling studies to aid in estimation of natural mortality and other factors affecting the tag return rate.
- Compare M and F estimates from acoustic tagging programs to conventional tagging programs.

Moderate

- Examine methods to estimate temporal variation in natural mortality.

Low

- Evaluate truncated matrices to reduce bias in years with no tag returns and covariate based tagging models to account for potential differences from size or sex or other covariates.

Life History, Biological, and Habitat Priorities

High

- Continue in-depth analysis of migrations, stock compositions, sex ratio, etc. using mark-recapture data.
- Continue evaluation of striped bass dietary needs and relation to health condition.
- Continue analysis to determine linkages between the *Mycobacteriosis* outbreak in Chesapeake Bay and sex ratio of Chesapeake spawning stock, Chesapeake juvenile production, and recruitment success into coastal fisheries.

Moderate

- Examine causes of different tag based survival estimates among programs estimating similar segments of the population.
- Continue to conduct research to determine limiting factors affecting recruitment and possible density implications.
- Conduct study to calculate the emigration rates from producer areas now that population levels are high and conduct multi-year study to determine inter-annual variation in emigration rates.

Additional Habitat Research Recommendations

- Passage facilities should be designed specifically for passing striped bass for optimum efficiency at passing this species.
- Conduct studies to determine whether passing migrating adults upstream earlier in the year in some rivers would increase striped bass production and larval survival, and opening downstream bypass facilities sooner would reduce mortality of early emigrants (both adult and early-hatched juveniles).
- All state and federal agencies responsible for reviewing impact statements and permit applications for projects or facilities proposed for striped bass spawning and nursery areas

shall ensure that those projects will have no or only minimal impact on local stocks, especially natal rivers of stocks considered depressed or undergoing restoration.

- Federal and state fishery management agencies should take steps to limit the introduction of compounds which are known to be accumulated in striped bass tissues and which pose a threat to human health or striped bass health.
- Every effort should be made to eliminate existing contaminants from striped bass habitats where a documented adverse impact occurs.
- Water quality criteria for striped bass spawning and nursery areas should be established, or existing criteria should be upgraded to levels that are sufficient to ensure successful striped bass reproduction.
- Each state should implement protection for the striped bass habitat within its jurisdiction to ensure the sustainability of that portion of the migratory stock. Such a program should include: inventory of historical habitats, identification of habitats presently used, specification of areas targeted for restoration, and imposition or encouragement of measures to retain or increase the quantity and quality of striped bass essential habitats.
- States in which striped bass spawning occurs should make every effort to declare striped bass spawning and nursery areas to be in need of special protection; such declaration should be accompanied by requirements of non-degradation of habitat quality, including minimization of non-point source runoff, prevention of significant increases in contaminant loadings, and prevention of the introduction of any new categories of contaminants into the area. For those agencies without water quality regulatory authority, protocols and schedules for providing input on water quality regulations to the responsible agency should be identified or created, to ensure that water quality needs of striped bass stocks are met.
- ASMFC should designate important habitats for striped bass spawning and nursery areas as HAPC.
- Each state should survey existing literature and data to determine the historical extent of striped bass occurrence and use within its jurisdiction. An assessment should be conducted of those areas not presently used for which restoration is feasible.

Management, Law Enforcement, and Socioeconomic Priorities

Moderate

- Examine the potential public health trade-offs between the continued reliance on the use of high minimum size limits (28 inches) on coastal recreational anglers and its long-term effects on enhanced PCB contamination among recreational stakeholders.