


*draft working paper for peer review only*



# Gulf of Maine winter flounder

## *2020 Assessment Update Report*

U.S. Department of Commerce  
National Oceanic and Atmospheric Administration  
National Marine Fisheries Service  
Northeast Fisheries Science Center  
Woods Hole, Massachusetts

Compiled August 2020

---

This assessment of the Gulf of Maine winter flounder (*Pseudopleuronectes americanus*) stock is a management track assessment of the existing 2017 area-swept operational assessment (NEFSC 2017). Based on the previous assessment the biomass status is unknown but overfishing was not occurring. This assessment updates commercial and recreational fishery catch data, research survey indices of abundance, and the area-swept estimates of 30+ cm biomass based on the fall NEFSC, MDMF, and MENH surveys.

**State of Stock:** Based on this updated assessment, the Gulf of Maine winter flounder (*Pseudopleuronectes americanus*) stock biomass status is unknown and overfishing is not occurring (Figures 1-2). Retrospective adjustments were not made to the model results. Biomass (30+ cm mt) in 2019 was estimated to be 2,862 mt (Figure 1). The 2019 30+ cm exploitation rate was estimated to be 0.052 which is 23% of the overfishing exploitation threshold proxy ( $E_{MSY}$  proxy = 0.23; Figure 2).

Table 1: Catch and status table for Gulf of Maine winter flounder. All weights are in (mt) and  $E_{Full}$  is the exploitation rate on 30+ cm fish. Biomass is estimated from survey area-swept for non-overlapping strata from three different fall surveys (MENH, MDMF, NEFSC) using an updated q estimate of 0.71 on the wing spread from the sweep study (Miller et al., 2020).

	2014	2015	2016	2017	2018	2019
	<i>Data</i>					
Recreational discards	5	5	11	5	2	2
Recreational landings	89	85	41	161	80	42
Commercial discards	5	2	3	3	3	4
Commercial landings	215	179	185	210	158	102
Catch for Assessment	315	271	241	378	244	150
	<i>Model Results</i>					
30+ cm Biomass	3,924	2,815	3,156	3,380	2,898	2,862
$E_{Full}$	0.08	0.096	0.076	0.112	0.084	0.052

Table 2: Comparison of reference points estimated in an earlier assessment and from the current assessment update. An  $E_{40\%}$  exploitation rate proxy was used for the overfishing threshold and was based on a length based yield per recruit model from the 2011 SARC 52 benchmark assessment.

	2017	2020
$E_{MSY}$ proxy	0.23	0.23
$B_{MSY}$	Unknown	Unknown
MSY (mt)	Unknown	Unknown
Overfishing	No	No
Overfished	Unknown	Unknown

**Projections:** Projections are not possible with area-swept based assessments. Catch advice was based on 75% of  $E_{40\%}$  (75%  $E_{MSY}$  proxy) using the terminal year fall area-swept estimate

assuming  $q=0.71$  on the wing spread which was updated using the average efficiency from 2009-2019 from the sweep experiment (Miller et al., 2020). Updated 2019 fall 30+ cm area-swept biomass (2,862 mt) implies an OFL of 658 mt based on the  $E_{MSY}$  proxy and a catch of 494 mt for 75% of the  $E_{MSY}$  proxy. Alternatively, using the average updated 2018 and 2019 fall 30+ cm area-swept biomass (2,880 mt) implies an OFL of 662 mt based on the  $E_{MSY}$  proxy and a catch of 497 mt for 75% of the  $E_{MSY}$  proxy.

### Special Comments:

- What are the most important sources of uncertainty in this stock assessment? Explain, and describe qualitatively how they affect the assessment results (such as estimates of biomass,  $F$ , recruitment, and population projections).

*The largest source of uncertainty with the direct estimates of stock biomass from survey area-swept estimates originates from the survey gear catchability ( $q$ ). Biomass and exploitation rate estimates are sensitive to the survey  $q$  assumption. However this 2020 update does incorporate the use of a re-estimated  $q$  through an average estimate of efficiency from 2009-2019 ( $q=0.71$ ) from the sweep study for the NEFSC survey. This updated  $q$  assumption (0.71) results in a higher estimate of 30+ biomass (2,862 mt) relative to the 2017 estimate  $q=0.87$  assumption (2,343 mt) from the updated fall surveys. Another major source of uncertainty with this method is that biomass based reference points cannot be determined and overfished status is unknown.*

- Does this assessment model have a retrospective pattern? If so, is the pattern minor, or major? (A major retrospective pattern occurs when the adjusted SSB or  $F_{Full}$  lies outside of the approximate joint confidence region for SSB and  $F_{Full}$ )

*The model used to determine status of this stock does not allow estimation of a retrospective pattern. An analytical stock assessment model does not exist for Gulf of Maine winter flounder. An analytical model was no longer used for stock status determination at SARC 52 (2011) due to concerns with a strong retrospective pattern. Models have difficulty with the apparent lack of a relationship between a large decrease in the catch with little change in the indices and age and/or size structure over time.*

- Based on this stock assessment, are population projections well determined or uncertain? If this stock is in a rebuilding plan, how do the projections compare to the rebuilding schedule?

*Population projections for Gulf of Maine winter flounder do not exist for area-swept assessments and stock biomass status is unknown. Catch advice from area-swept estimates tend to vary with interannual variability in the surveys. Consideration should be given to using multiple surveys to stabilize the biomass estimates and catch advice.*

- Describe any changes that were made to the current stock assessment, beyond incorporating additional years of data and the effect these changes had on the assessment and stock status.

*The assumption on  $q$  changed from 0.87 to 0.71 using information from the updated sweep experiment (Miller et al., 2020) and incorporation of new survey data were made to this Gulf of Maine winter flounder management track assessment. The new MRIP calibrated catch time series was also updated in this assessment. In addition there were some changes with updated commercial landings data with the switch to using Stockeff data which are mostly due to the changes in the proration with regards to unknown areas from Massachusetts*

*state landings of winter flounder. However, Changes in total removals will not affect the biomass or catch advice and total removals still remain far below the overfishing definition.*

- If the stock status has changed a lot since the previous assessment, explain why this occurred.

*The overfishing status of Gulf of Maine winter flounder has not changed.*

- Provide qualitative statements describing the condition of the stock that relate to stock status.

*The Gulf of Maine winter flounder has relatively flat survey indices with little change in the size structure over time. There have been large declines in the commercial and recreational removals since the 1980s. However, this large decline over the time series does not appear to have resulted in a response in the stock's size structure within the catch and surveys nor has it resulted in a change in the survey indices of abundance.*

- Indicate what data or studies are currently lacking and which would be needed most to improve this stock assessment in the future.

*Direct area-swept assessments could be improved with additional studies on state survey gear efficiency. Quantifying the degree of herding between the doors and escapement under the footrope and/or above the headrope for state surveys is needed to improve the area-swept biomass estimates. Studies quantifying winter flounder abundance and distribution among habitat types and within estuaries could improve the biomass estimate.*

- Are there other important issues?

*The general lack of a response in survey indices and age/size structure are the primary sources of concern with catches remaining far below the overfishing level.*

#### **References:**

Northeast Fisheries Science Center. 2017. Operational Assessment of 19 Northeast Groundfish Stocks, Updated Through 2016. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 17-17; Commer, Northeast Fish Sci Cent Ref Doc. 17-17; 259 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026.  
<https://repository.library.noaa.gov/view/noaa/16091>

Northeast Fisheries Science Center. 2011. 52<sup>nd</sup> Northeast Regional Stock Assessment Workshop (52<sup>nd</sup> SAW) Assessment Report. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 11-17; 962 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026. [CRD11-17](#)




Figure 1: Trends in 30+ cm area-swept biomass of Gulf of Maine winter flounder between 2009 and 2019 from the current assessment based on the fall (MENH, MDMF, NEFSC) surveys.




Figure 2: Trends in the exploitation rates ( $E_{Full}$ ) of Gulf of Maine winter flounder between 2009 and 2019 from the current assessment and the corresponding  $F_{Threshold}$  ( $E_{MSY}$  proxy=0.23; horizontal dashed line).




Figure 3: Total catch of Gulf of Maine winter flounder between 2009 and 2019 by fleet (commercial and recreational) and disposition (landings and discards). A 15% mortality rate is assumed on recreational discards and a 50% mortality rate on commercial discards.




Figure 4: Indices of biomass for the Gulf of Maine winter flounder between 1978 and 2019 for the Northeast Fisheries Science Center (NEFSC), Massachusetts Division of Marine Fisheries (MDMF), and the Maine New Hampshire (MENH) spring and fall bottom trawl (strata 1-3) surveys. NEFSC indices are calculated with gear and vessel conversion factors where appropriate. The approximate 90% lognormal confidence intervals are shown.